# An Overview on Understanding the Basic Concept of Fish Diseases in Aquaculture

ALFRED, O.<sup>1</sup>, SHAAHU, A.<sup>2</sup>, ORBAN, D. A.<sup>3</sup>, EGWENOMHE, M.<sup>4</sup>

<sup>1, 4</sup> Department of Fisheries, University of Benin, Nigeria.
 <sup>2</sup> Soybean Research Programme, National Cereals Research Institute, Badeggi, Nigeria.
 <sup>3</sup> Department of Agriculture, College of Education Katsina-Ala, Benue State, Nigeria.

Abstract- Fish health management is concerned with the care and proper measures in managing the healthy living of fish in aquaculture. Disease issues are of great concern in aquaculture production. This is because disease is considered one of the important factors to decrease in fish production, both in farming system and in wild condition. Production costs are increased through investment lost in dead cultured animals, cost of treatment, and decreased quality and quantity of yields. This study became necessary as there were various efforts to increase fish supply through aquaculture production to bridge the gap between the declining fish supply from capture fisheries and the increasing food fish demand. This study reviews fish disease in aquaculture and is focused on the nature, types, causes and significance of fish diseases in aquaculture production to provide hands-on information on diseases and health management in aquaculture production, and create relevant awareness on cultured fisheries management and practices.

Indexed Terms- Aquaculture Development; Disease; Infectious; Non-infectious; Economic Loss;

# I. INTRODUCTION

Aquaculture is the fastest growing industry around the world and Nigeria is no exception. Aquaculture production, especially pond aquaculture, is depends upon sources of obtaining increased fish production in order to supply and feed the ever-increasing population of the world (FAO 2010). New technological advances and increased demands for fish as a source of animal protein are the main reasons for the industry's growth. Because of expansion of the industry, the culture methods have become more intensive for producing higher yields (Rico *et al.*,

2012). Recently, the sector attracted great attention and it is growing rapidly through the development of aquaculture (Kubecka et al., 2016). The advent and anticipation of aquaculture is to enhance food production; more income generation and good health for the people. However, like other farming sectors, disease is a substantial source of constraint to aquaculture development and sustainability; from both social and economic points of view (Blanco et al., 2000; Allsopp et al., 2008 and Bostock et al., 2010). Huge loss of production in aquaculture is occurring because of many reasons. Among these causes, a disease is the most serious constraint that causes damage to the livelihood of farmers, loss of job, reduced incomes, and food insecurity. Disease outbreaks have long been recognised as a significant constraint to aquaculture production and economic viability (Sarig 1971 and Bondad-Reantaso et al., 2005). Studies showed that almost fifty percent of production loss is because of diseases which are more severe in developing countries. This is because ninety percent of the aquaculture firm is in the developing world. The annual loss of revenues because of disease reaches up to 6 billion dollars (Leung and Bates, 2013, Hien et al., 2020). Disease is a condition in living organisms in which normal physiological functions are being impaired due to alteration in the body systems and typically manifested by distinguishing signs and symptoms (pathological symptoms) (Idowu et al., 2016). Disease is one of the major constraints to aquaculture and limiting factor for economic and social developments in different countries of the world (Bagun et al., 2013). Production costs are increased through investment lost in dead cultured animals, cost of treatment, and decreased quality and quantity of vields (Francis 2005). Disease is considered one of the important factors to decrease in fish production, both in farming system and in wild condition. Large-scale mortality of fish often occurs in ponds due to

environmental stress followed by parasitic invasion and bacterial, fungal, protozoan and monogenean infections (Hossain *et al.*, 2011).

# II. NATURE AND TYPES OF FISH DISEASE

Healthy fish have adequate resistance against diseases; they can adapt to reasonable environmental changes and in turn resist diseases (AFCD 2009 and Govind et al., 2012). Nevertheless, diseases occur in fish, but before an active fish disease is developed in a culture system pathologically linked factors are involved. These include; Presence of environmental pathogens, Low resistance of the fish and Unfavorable water environment (Idowu et al., 2017 and Hien et al., 2020). Development of an active disease in fish results from the effect of the association among these (pathologically linked) factors. In a pond, when the pathogen load increases due to external factors (environmental problems such as poor water supply, or other stressors), above what the natural resistance of the fish can cope with, fish become vulnerable to pathogenic infections and diseases. Also, external factors may cause drastic changes in water quality and lower fish resistance. When these happen, fish become susceptible to diseases, even the risks of fish kill are heightened (Idowu et al., 2017).

# III. FORMS OF FISH DISEASES

There are two broad forms of diseases affecting fish. Fish diseases may be subdivided into:

- Infectious diseases
- Non-infection diseases

# 1) Infectious Diseases

This is also referred to as biotic disease. Infectious diseases are caused by living factors – pathogenic organisms (viruses, bacteria, fungi or parasites) present in the aquatic environment or carried by other fish. Fish become vulnerable to pathogenic infections when there are stressors (environmental abnormalities, water quality deterioration, unbalanced nutrition, or bodily injuries) which weaken fish natural resistance (immune system). Infectious diseases pose a unique problem of diagnosis (Govind et al., 2012). The emergence of infectious diseases is usually triggered by ecological changes, often associated with human interventions, such as transfer of organisms,

environmental degradation, agricultural practices or technology (Jones *et al.*, 2008). Infections can occur internally and externally affecting tissues, organs and other fish body parts. They are mostly contagious diseases and some types of treatments may be necessary to control the disease outbreaks (Idowu *et al.*, 2017).

- The Common Examples of Infectious Diseases Are Discussed Below.
- i. Parasitic Diseases

Parasites are organisms that live in (endo) or on (ecto) another organism (called host) of different species upon which it depends on nourishment at the expense of their host. They can be found in virtually all living things and in normal natural conditions exist side by side with their host. Most parasite species rarely cause problems in the natural environment but in aquaculture, parasites often cause serious outbreaks of disease (Roberts 2012). They play an important role in determining the productivity, sustainability, and economic viability of aquaculture. Parasite infections cause serious socioeconomic, ecological, and welfare consequences in global finfish aquaculture (Menezes et al., 1990, Barber 2007, Shinn et al., 2015). Fish parasites infest the gills, skin, gut, or as a grub-like worms in fish muscle tissue causing irritation, impaired function, weight loss, and eventually fish kill (Francis 2005). Parasites of fish are not strictly pathogens, but they make a convenient classification as infectious disease of fish (Paperna, 1996 and Fiala et al., 2015), as they are often accompanied or followed by secondary bacterial or fungal infection.

- Examples of Parasitic Disease Are Given Below with Their Various Causative Organisms.
- i) Protozoan Diseases
- Ichthyophthiriasis (Ichthyophthirius multifiliis)
- Ichthyobodosis (Ichthyobodo necatrix)
- Chilodonelliasis (Chilodonella hexasticha)
- Amoebic gill disease (Neoparamoeba perurans)
- ii) Metazoan Diseases
- Whirling Disease (*Myxobolus cerebralis*)
- Diplostomiasis (Diplostomum spathaceum)

- Proliferative kidney disease (*Tetracapsuloides* bryosalmonae)
- Lernaeosis (Lernaea cyprinacea)

iii) Zoonotic Diseases

- Capillariasis (*Capillaria philippinensis*)
- Diphyllobothriasis (*Diphyllobothrium latum*)

Source: (Ali and Faruk, 2018, Idowu et al., 2017).

#### ii. Bacterial Diseases

Bacteria are usual component of aquatic environment. They become problem when fish are exposed to stressors (Omojowo and Sogbesan, 2003). Bacterial diseases are responsible for heavy mortality in both wild and cultured fish. The actual role of these microorganisms may vary from that of a primary pathogen to that of an opportunist invader of a host rendered moribund by some other disease process (Richards & Roberts, 1978 and Abowei & Brivai, 2011). The specific susceptibility of aquatic animals to various bacteria can be observed only in the species that are cultured, or atleast kept in captivity for a while. Bacterial infections are considered the major cause of mortality in aquaculture (Govind et al., 2012). Bacteria are very common in the aquatic environment. Most bacterial disease agents are part of the normal flora of the water. They cause disease only when the fish are stressed due to poor environmental conditions, inadequate diet and poor husbandry techniques (Tendencia, & Lavilla-Pitogo, 2004). Bacteria are important pathogens for both cultivated and wild fish, and are responsible for serious economic losses. Some bacteria cause only surface diseases as skin or gill infections, especially flexibacteria, but some inflict systemic disease (Inglis et al., 2001). They can infect a single fish and multiply rapidly to cause a substantial fish kill in a few days or weeks. Bacterial diseases are often internal infections and usually require treatment (with antibiotics added to feeds or water). However, bacterial diseases can also be external which may result from rough handling or effects of parasitic infection (Idowu et al., 2017). Bacterial diseases are often internal infections

- Examples of Some Important Bacterial Diseases in Aquaculture with their causative organisms
- i) Columnaris Disease (Flavobacterium columnare)

- ii) Enteric Septicemia Disease (Edwardsiella ictaluri)
- iii) Coldwater (Peduncle) Disease (*Flavobacterium* psychrophilum)
- iv) Bacterial kidney disease (*Renibacterium* salmoninarum)
- v) Franciselliosis (Francisella noatunensis)
- vi) Furunculosis (Aeromonas salmonicida)
- vii)Myxobacteriosis (Fiexibacter maritimu)
- viii) Edwardsiliosis (Edwardsiella tarda)
- ix) Fin rot (Flexibacter maritimus)
- x) Vibriosis (Vibrio sp)

Source: (Tendencia, & Lavilla-Pitogo, 2004 and Pekala & Agnieszka,2018).

#### iii. Viral Diseases

Viruses are very small infectious agents that multiply only within the living cells of an animal or plant host. Because they are so small, viruses are often difficult to detect. Viruses occur in particles and they are obligate pathogens. They depend on the synthesizing structure of the host cells for replication. Viral diseases of fish are difficult to diagnose and control with medications (Francis, 2005). The viruses affecting the fishes grow at lower temperatures and have specific tolerance to temperature. Most of the viral diseases affecting the fishes are geographically limited. In most of the viral infections, it causes high mortality in fishlings but adults become carrier to these viruses (Ganguly, 2017). Fish are susceptible to many viral pathogens, some of which are readily identifiable, whereas others remain obscure. Identification is not always possible owing to lack of resources or inability of the virus to grow in available fish cell lines. In many instances viral infection may remain latent until adverse environmental conditions, such as poor water quality, overcrowding, or rough handling, trigger the onset of disease (Bernoth & Crane, 1995). Some viral infections are serious diseases of groupers causing heavy mortalities. In most cases, larval stages are the most susceptible stage. With the carnivorous nature of groupers, they can readily ingest viral pathogens from live fish food or trash fish that carry the viral pathogens. Moreover, viruses are able to effect vertical transmission from broodstocks that are likely carriers of the virus (Lio-Po et al., 2004). Viruses are probably the cause of a wide spectrum of fish diseases. Although relatively few virus diseases of fish are known today, some of the diseases of unknown

# © DEC 2020 | IRE Journals | Volume 4 Issue 6 | ISSN: 2456-8880

etiology, as well as some diseases presently accepted as due to bacteria, protozoa, fungi or nutritional deficiencies, possibly will be recognized eventually as virus diseases (Stanely, 1954 and Wise *et al.*, 2004).

- Examples of Some Important Viral Diseases in Aquaculture with their causative organisms
- i) Lymphocystis disease (Lymphocystivirus)
- ii) Infectious Pancreatic Necrosis (Aquabirnavirus)
- iii) Viral Haemhorragic Septicaemia (Novirhabdovirus)
- iv) Infectious Haematopoietic Necrosis (Novirhabdovirus)
- v) Nodavirosis (Betanodavirus)
- vi) Goldfish Iridovirus
- vii)Snakehead Rhabdovirus
- viii) Bluegill Hepatic Necrosis Reovirus
- ix) Grass Carp Reovirus
- x) Channel Catfish Virus Disease

Source: (Barja, 2004 and WOLF, 1988)

#### iv. Fungal Diseases

Fungal diseases of fish have become increasingly important over the past 20 years. Fungal infections in fish generally are considered secondary to some other factor or pathogen, a consequence of water quality problems, poor condition, trauma (rough handling or aggression), bacterial disease, or parasites. Fungi, however, can cause disease under a variety of other circumstances. Some may be more aggressive and play a more primary role. Fungi can be external or internal and systemic (Yanong, 2003). The fungal diseases occur in brood stock as well as all life stages of fish and eggs. Fungal infection causes low yield of fry and low production in fish culture (Kwanprasert et al., 2007). Fungal infections (fungal infections are called mycoses) are among the most common diseases seen in temperate fish because fungal spores are found in all fish ponds and create problems in stressed fish. Poor water quality can also lead to an increase in fungal infections in an otherwise healthy fish population. Most fungal infections invaded on external tissues and only few fungal infections affect the internal organs of fish (Verma, 2008).

• Examples of Some Important Fungal Infection in Aquaculture with their causative organisms

- ii) Dermocystidiosis (Dermocystidium marinum)
- iii) Ichthyophonosis (Ichthyophonus hoferican)
- iv) Exophialiasis (Exophiala salmonis)
- v) Branchiomycosis (Branchiomyces sanguinis)
- vi) Aspergillomycosis (Aspergillus spp)

Source: (patel et al., 2018)

#### 2) NON-INFECTIOUS DISEASES

Non-infectious diseases (also referred to as systemic diseases) are caused by non-living/abiotic factors. Compared with infectious diseases and disorders, few non-infectious diseases and disorders in cultured fish have severe biologic or economic impact. Culture practices, however, often establish environments that promote infectious disease by weakening the immune response or by promoting conditions that favor infectious agents. Generally non-infectious diseases and disorders result from adverse physical, chemical and biological factors and from poor nutrition. Adverse factors can be changes in temperature, pH, and dissolved gases, suspended solids, endogenous toxins, anthropogenic toxicants, solar radiation and physical damage such as from predation (Hawkins, 2002). The diseases are either congenital (such as genetic anomalies or neo plastic conditions) or iatrogenic (induced by external conditions such as environmental or nutritional problems) (Idowu et al., 2017).

Non-infectious diseases can be broadly categorized as;

- Environmental
- Nutritional
- Genetic.

#### i. Environmental Diseases

Environmental diseases are the most important among non-infectious commercial the diseases in aquaculture. Environmental diseases result from inadequacies in the physical and chemical characteristics of the pond water. These include low dissolved oxygen, high ammonia, high nitrite or natural or man-made toxins in the aquatic environment (Francis, 2005). Environmental problems include depletion of dissolved oxygen, extremes in pH, high ammonia, high nitrite, natural or man-made toxins, or mechanical trauma (caused by rough handling, overcrowding, low water levels, or predation). Proper

i) Saprolegniasis (Saprolegnia parasitica)

water quality management and handling are necessary in preventing most environmental problems (Idowu *et al.*, 2017).

- Examples of Some Frequently Occurring Environmental Diseases in Aquaculture
- i) Gas Bubble Disease
- ii) Swimbladder Stress Syndrome
- iii) Asphyxiation/Hypoxia
- iv) Sunburn Disease
- v) Brown Blood Disease
- vi) Acidosis and Alkalosis Disease

Source: (Erazo-Pagador, 2001., Erazo-Pagador, & Cruz-Lacierda, 2004)

#### ii. Nutritional Diseases

Nutritional diseases are those which the results in due to excess nutrient or nutritional deficiency in fish than the normal requirements (Idowu et al., 2017). Nutritional diseases of fish may develop as a result of deficiency (undernutrition), excess (overnutrition), or imbalance (malnutrition) of nutrients present in their food. The disease usually develops gradually because animals have body reserves that make up for nutritional deficiency up to a certain extent (Amar & Lavilla-Pitogo, 2004). Symptoms appear gradually when one or more components in the diet drop below the critical level of the body reserves. Most nutritional diseases are difficult to diagnose because of their chronic nature, with the condition only manifesting over a long period of time (Abowei et al., 2011). When there is too much food, the excess that is converted to fat and deposited in fish tissues and organs, may severely affect physiological functions of the fish (Amar & Lavilla-Pitogo, 2004). Nutritional diseases arising from dietary imbalances, continues to cause problems to fish in cultured condition (Aly, 2013). Diets that are inadequate with respect to protein, amino acid, essential fatty acids, vitamins and minerals lead to gross malnutrition and high disease susceptibility. Proper feeding of a nutritious diet is important for growth and prevention of nutritional deficiencies, and to cope with a variety of diseasecausing agents (Lall, 2000). Signs of nutritional disease can also be marked by secondary disease condition due to pathogens (Blanco et al., 2000).

- Examples of Some Nutritional Disease in Aquaculture
- i) Lipodosis
- ii) Fish scurvy
- iii) Broken Back Syndrome
- iv) Steatitis and white fat disease
- v) Avitaminosis

Source: (Shoaibe and Mohammed, 2018)

#### iii. Genetic abnormalities

Abnormalities in fish include any changes that are obvious by looking at the outside (external) or inside (internal) of the fish. This can be discussed as Neo plastic And Genetic Anomalies. A neoplasm is an abnormal growth of cells, also known as a tumor. Neo plastic fish disease refers to abnormal growth in any of the organs with resultant loss of structural and functional ability of the affected organ. The resultant growth may be lethal or mildly pathologic (Abowei et al., 2011). The hereditary factors involved in diseases and abnormalities are generally complex and difficult to detect, but the evidence for their existence is undeniable (Bernard and Michel. 1990). Malformations occurring in cultured fish or induced following hybridizations are of minimal significance; however, it is of importance to bring in unrelated fish for use as brood stock every few years to minimize inbreeding (Francis, 2005). This way, genetic abnormalities can be controlled.

# IV. THE SIGNIFICANCE OF FISH DISEASE TO AQUACULTURE

Aquaculture industry plays a vital role in many countries in the world especially developing countries. Fish disease is a substantial source of monetary loss to aquaculturists. Production costs are increased by fish disease outbreaks because of the investment lost in dead fish, cost of treatment, and decreased growth during convalescence (Francis, 2005). Fish disease impedes both economic and social development in many ways: directly, through production losses and increased operational costs and indirectly, through cost to society (social, welfare and environmental), adjustment in market shares and increase in price due to lower supply. Fish diseases affect fish survival and growth rates resulting to poor yield (both in quality and quantity), the livelihood of people involved in the culture production and the community in which they occur through reduced food availability, loss of earnings/employments and recreation, apprehension of healthy environment, consumption and handling of sick fish. Invariably, fish culturists and consumers suffer hefty socio-economic losses (Francis, 2005; Bondad *et al.*, 2005; Brun *et al.*, 2009; Oladele *et al.*, 2011; Ali *et al.*, 2014).

World Bank in 2006 reported global loss of about US \$3billion per year to aquaculture production and trade due to disease (Subasinghe *et al.*, 2001, Brun *et al.*, 2009). Significance of fish diseases can also be in form of investment in disease research and control, along with health management programmes (Bondad *et al.*, 2005). Reported on the financial implications incurred from fish disease control programmes and fish disease legislation by various countries. Moreover, discovery of pathogens and unapproved drugs in aquaculture products has also continued to generate issues including rejection of products, enforcement actions against involved trade parties (country, industry, importer, etc.), serious trade disruption and heavy financial losses (Karunasagar, 2012).

#### CONCLUSION

Fish health management is an important area of aquaculture that needs to be taken seriously. Poor health management weakens the immune system of fish, thereby predisposes fish to disease and infection. Disease has become a primary constraint to sustainable aquaculture production and products trade, thereby affect the socioeconomic status of most developing countries. Though enormous loss is incurred due to diseases, less attention is given for disease management and a very few well trained persons are available in this field (Subasinghe et al., 2001 and Uddin et al., 2017). Diseases have brought about major setback in the development of the sector. Disease reduces supply of aquaculture products, increases cost of production, loss of income to the farmer, reduce quality of fish produced and many more effects. The rural farmers are still the most involving in aquaculture production (Faruk et al., 2004). Most of them are with little or no knowledge of aquaculture health management and with inadequate opportunities to improve management skills and respond effectively to disease problems. More

enlightenment on the dangers and effects of disease should be preached to farmers across the globe. This way, the damage that is usually impaired by diseases on aquaculture will be reduced if not totally eradicated. The nature and types of disease is also and help or a guide as to know how effectively to tackle the rampant disease problem. Most importantly, the infectious diseases which are causes by several pathogens, this has to be quickly tackled because the danger of this disease is that is spreads rapidly among's the stock and can totally eliminate the full stock in a farm. Both the infectious and non-infectious disease is sometimes due to the negligence of farmers. Generally, prevention of this disease is far better than the control or treatment.

#### REFERENCES

- [1] Abowei, J. & Briyai, O. (2011). A Review of Some Bacteria Diseases in Africa Culture Fisheries. Asian Journal of Medical Sciences. 3.
- [2] Abowei JFN, Briyai OF, Bassey SE (2011) A review of some viral, neoplastic, environmental and nutritional diseases of African fish. British Journal of Pharmacology and Toxicology 2(5): 227-235.
- [3] Agriculture, Fisheries and Conservation Department (2009) Prevention and treatment of fish diseases. Good Aquaculture Practices Series. Aquaculture Fisheries Division, AFCD, Hong Kong (4): 32.
- [4] Ali, Md and Faruk, Md. (2018). Fish Parasite: Infectious Diseases Associated with Fish Parasite.
- [5] Omojowo FS, Sogbesan OA (2003) Fish losses due to bacterial flora and infections of fishes in kainji lake Fish Diseases, Nigeria: A review. Nigerian Veterinary Journal 24: 41-47.
- [6] Ali MH, Chowdhury FS, Ashrafuzzaman M, Chowdhury MA, Ul-Haque MR, et al. (2014) Identification, pathogenicity, antibiotic and herbal sensitivity of Edwardsiella tarda causing fish disease in Bangladesh. Current Research in Microbiology and Biotechnology 2(1): 292-297.

- [7] Allsopp M, Johnston P, Santillo D (2008) Challenging the aquaculture industry on sustainability: Technical overview. Greenpeace Research Laboratories Technical Note 01: 1-59.
- [8] Aly Salah Mesalhy. "A review of fish diseases in the Egyptian aquaculture sector: Working report". (2013).
- [9] Amar, E. C., & Lavilla-Pitogo, C. R. (2004). Nutritional diseases. In K. Nagasawa & E. R. Cruz-Lacierda (Eds.), Diseases of cultured groupers (pp. 59-66). Tigbauan, Iloilo, Philippines: Aquaculture Department, Southeast Asian Fisheries Development Center.
- [10] Bagum N., Monir M.S. and Khan M.H., (2013). Presents status of fish diseases and economic losses due to incidence of disease in rural freshwater aquaculture of Bangladesh. *J Innov Dev Strategy*, 7(3):48-53.
- [11] Barber, I. 2007. Parasites, behaviour and welfare in fish. Appl. Anim. Behav. Sci. 104: 251–264
- [12] Barja J.L. Report about fish viral diseases.In : Alvarez-Pellitero P. (ed.), Barja J.L. (ed.),
- [13] Basurco B.(ed.), Berth e F. (ed.), Toranzo A.E. (ed.). Mediterranean aquaculture diagnostic laboratories. Zaragoza: CIHEAM, 2004. p. 9 1 1 02 (Option s Méditerranéennes: Série B. Etudes et Recherches; n. 49)
- [14] Bernard Chevassus and Michel Dorson, Genetics of resistance to disease in fishes, *Aquaculture*, 10.1016/0044-8486(90)90009-C, **85**, 1-4, (83-107), (1990).
- [15] Bernoth E.-M., Crane M.S.J. Viral diseases of aquarium fish (1995) *Seminars in Avian and Exotic Pet Medicine*, 4 (2), pp. 103-110.
- [16] Blanco MM, Gibello A, Fernández-Garayzábal JF (2000) Influence of fish health management: Bases, procedures and economic implications. Global quality assessment in Mediterranean aquaculture Zaragoza CIHEAM 51: 45-49.
- [17] Bondad Reantaso MG, Subasinghe RP, Arthur J R, Ogawa K, Chinabut S (2005) Disease and health management in Asian aquaculture. Veterinary Parasitology VETPAR 3314: 24.

- [18] Bostock, J., McAndrew, B., Richards, R., Jauncey, K., Telfer, T., Lorenzen, K., Little, D., Ross, L., Handisyde, N., Gatward, I. and Corner, R., 2010. Aquaculture: global status and trends. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1554), pp.2897-2912.
- [19] Brun E, Rodgers C, Georgiadis M, Bjorndal T (2009) Economic impact of disease and biosecurity measures. International Biosecurity Conference of National Veterinary Institute on Economic Impact of Disease and Biosecurity Measures 17-18.
- [20] Economic costs of protistan and metazoan parasites to global mariculture. Parasitology 142: 196–270
- [21] Erazo-Pagador, G. E. (2001). In G. D. Lio-Po, C.
  R. Lavilla, & E. R. Cruz-Lacierda (Eds.), Health management in aquaculture (pp. 75-81).
  Tigbauan, Iloilo, Philippines: Aquaculture Department, Southeast Asian Fisheries Development Center.
- [22] Erazo-Pagador, G. E., & Cruz-Lacierda, E. R. (2004). Environmental diseases. In K. Nagasawa & E. R. Cruz-Lacierda (Eds.), Diseases of cultured groupers (pp. 67–71). Tigbauan, Iloilo, Philippines: Aquaculture Department, Southeast Asian Fisheries Development Center.
- [23] FAO, 2010. State of World Aquacultutre-Fisheries Department. FAO Fisheries Technical Paper 500,21-26.
- [24] Faruk MAR, Sarker MMR, Alam MJ, Kabir M B (2004) Economic loss from fish diseases on rural freshwater aquaculture of Bangladesh. Pakistan Journal of Biological Sciences 7(12): 2086-2091.
- [25] Fiala, I., Bartošová-Sojková, P. and Whipps, C.M., 2015. Classification and phylogenetics of Myxozoa. In *Myxozoan evolution, ecology and development* (pp. 85-110). Springer, Cham.
- [26] Francis Floyd R (2005) Introduction to fish health management. IFAS Extension. Fisheries and Aquatic Sciences Department University of Florida US CIR 921: 1-4.

- [27] Ganguly, S., Viral Diseases Infecting Finfishes and Ornamental Fishes: A Review of Relevance to Sustainable Aquaculture, Int. J. Pure App. Biosci.5(1):282-284(2017). doi: http://dx.doi.org/10.18782/2320-7051.2455
- [28] Govind P, Madhuri S, Shrivastav AB Sahni YP (2012) Overview of the treatment and control of common fish diseases. International Research Journal of Pharmacy 3(7): 123-127.
- [29] Hawkins, W. E., J W. Fournie, AND N. Chansue. NON-INFECTIOUS DISORDERS OF WARMWATER FISHES. 2002, P.T.K. Woo, D.W. Bruno, S.L.H. Lim (ed.), Diseases of Finfish in Cage Culture. CAB International, Wallingford, Uk, , 283-304.
- [30] Hien V. D., Seyed H. H., Einar R., Maria A. E, Maryam D., Mahmoud A. O. Dawood & Caterina F. (2020) Host-Associated Probiotics: A Key Factor in Sustainable Aquaculture, Reviews in Fisheries Science & Aquaculture, 28:1, 16-42, DOI: 10.1080/23308249.2019.1643288
- [31] Hossain, M.K., Islam, K.T., Hossain, M.D. and Rahman M.H., (2011). Environmental impact assessment of fish diseases on fish production. J. Sci. Foundation 9(1&2): 125-131.
- [32] Idowu TA, Onyia LU, Kefas M (2016) Fish diseases and health management. In: Contextual aquaculture and fisheries digest. Maiden Edition Paraclete Publisher 155-171.
- [33] Idowu TA, HA Adedeji and OA Sogbesan (2017) Fish Disease and Health Management in Aquaculture Production. Int J Environ & Agri Sci 1: 002.
- [34] Inglis V., Roberts R.J., Bromage N.R. 2001.Bacterial Diseases of Fish. Iowa State University, Ames. pp. 1 –59, 122 –156.
- [35] Jones K. E., Patel N. G., Levy M. A., Storeygard A., Balk D., Gittleman J. L., Daszak P.2008Global trends in emerging infectious diseases. Nature 451, 990–993
- [36] Karunasagar I (2012). Public health and trade impact of antimicrobial use in aquaculture. In: MG Bondad-Reantaso, JR Arthur and RP Subasinghe, eds Improving biosecurity through

prudent and responsible use of veterinary medicines in aquatic food production 1-9. FAO Fisheries and Aquaculture Technical Paper No. 547 Rome FAO 207.

- [37] Kubecka J., Boukal D. S., Cech M. et al., "Ecology and ecological quality of fish in lakes and reservoirs," *Fisheries Research*, vol.173, pp. 1–3, 2016.
- [38] Kwanprasert P, Hangavant C, Kitancharoen N (2007) Characteristics of Achylabisexualis Isolated from Eggs of Nile Tilapia (OreochromisniloticusLinn.). KKU Res J 12: 195-202.
- [39] Lall Santosh P. "Nutrition and health of fish". Avances en Nutrición Acuícola V. Memorias del V Simposium Internacional de Nutrición Acuícola (2000): 19-22.
- [40] Leung T. L. F. and Bates A. E., "More rapid and severe disease outbreaks for aquaculture at the tropics: Implications for food security," *Journal* of Applied Ecology, vol. 50, no. 1, pp. 215–222, 2013.
- [41] Lio-Po, G. D., & de la Peña, L. D. (2004). Viral diseases. In K. Nagasawa & E.R. Cruz-Lacierda (Eds.), Diseases of cultured groupers (pp. 3-18). Tigbauan, Iloilo, hilippines:Aquaculture Department, Southeast Asian Fisheries Development Center.
- [42] Menezes J., Ramos M.A., Pereira T.G. and Moreira da Silva, A. 1990. Rainbow trout culture failure in a small lake as a result of massive parasitosis related to careless introductions. Aquaculture 89: 123–126
- [43] Oladele OO, Olufemi BE, Oladosu GA, Ajayi OL, Adediji AA, et al. (2011) Arborescent organ necrosis syndrome in cat fish Clarias gariepinus (Burchell): a case report. Journal of Fish Diseases 34(10): 801-804.
- [44] Paperna I(1996) Parasites, infection and diseases of fishes in Africa-An update. CIFA Technical Paper No 31. Rome FAO: 220.
- [45] Patel A.S., Patel S.J., Bariya A.R., Pata B.A. and Ghodasara S.N. Fungal Diseases of Fish: A Review. Vet Sci Res 2018, 3(3): 000164.

- [46] Pekala, Agnieszka. (2018). Contemporary Threats of Bacterial Infections in Freshwater Fish. Journal of Veterinary Research. 62. 261-267. 10.2478/jvetres-2018-0037.
- [47] Richards, R.H. & Roberts, R.J., 1978. The bacteriology of teleosts. In: Roberts, R.J. (ed.) Fish Pathology. Bailliere, Tindall, London. pp. 183–204.
- [48] Rico, K. Satapornvanit, M. M.Haque et al., "Use of chemicals and biological products inAsian aquaculture and their potential environmental risks: A critical review," *Reviews in Aquaculture*, vol. 4, no. 2, pp. 75–93, 2012.
- [49] Roberts, R.J. 2012. The parasitology of teleosts.In: Fish Pathology, Fourth Edition. Blackwell Publishing Ltd
- [50] Sarig, S. (1971). Diseases of Fishes. Book 3: The Prevention and Treatment of Diseases of Warmwater Fishes under Subtropical Conditions, with Special Emphasis on Intensive Fish Farming. TFH Publications, Inc. Ltd., London.
- [51] Shinn, A.P., Pratoomyot, J., Bron, J.E., Paladini, G., Brooker, E.E. and Brooker, A.J. 2015.
- [52] Shoaibe Hossain Talukder Shefat and Mohammed Abdul Karim. "Nutritional Diseases of Fish in Aquaculture and Their Management: A Review". Acta Scientific Pharmaceutical Sciences 2.12 (2018): 50-58.
- [53] Stanely W. Watson (1954) Virus Diseases of Fish, Transactions of the American Fisheries Society, 83:1, 331-341, DOI: 10.1577/1548-8659(1953)83[331: VDOF]2.0.CO;2
- [54] Subasinghe RP, Bondad Reantaso MG, McGladdery SE (2001) Aquaculture development, health and wealth. In: Subasinghe RP, Bueno PB, Phillips MJ, Hough C, McGladdery SE, Arthur JR (Eds.). Technical Proceedings of the Conference on Aquaculture in the Third Millenium, Bangkok, Thailand, 20-25. February 2000. NACA, Bangkok and FAO, Rome: 471.
- [55] Tendencia, E. A., & Lavilla-Pitogo, C. R. (2004). Bacterial diseases. In K. Nagasawa & E. R. Cruz-

Lacierda (Eds.), Diseases of cultured groupers (pp. 19-28). Tigbauan, Iloilo, Philippines: Aquaculture Department, Southeast Asian Fisheries Development Center.

- [56] Verma V (2008) Fungus disease in fish, diagnosis and treatment. Veterinary World1(2): 62.
- [57] Wise, D.J., Camus, A.C., Schwedler, T.E. and Terhune, J.S., 2004. 15 Health management. In *Developments in Aquaculture and Fisheries Science* (Vol. 34, pp. 444-503). Elsevier.
- [58] WOLF, K. (1988). Fish Viruses and Fish Viral Diseases. ITHACA; LONDON: Cornell University Press. doi:10.7591/j. ctvr7f4z5
- [59] Yanong, Roy. (2003). Fungal diseases of fish. The veterinary clinics of North America. Exotic animal practice. 6. 377-400. 10.1016/S1094-9194(03)00005-7.