

Fekete-Szegö bounds for a Subclass of Starlike and Inverse Starlike Analytic Functions

KONIKA

Student, GSSDGS Khalsa College, Patiala

Abstract- Here we describe some classes of analytic functions and its subclasses by which we will be obtaining sharp upper bounds of the functional $|a_3 - \mu a_2^2|$ for the analytic function $f(z) = z + \sum_{n=2}^{\infty} a_n z^n, |z| < 1$ belonging to these classes and subclasses.

Indexed Terms- Univalent functions, Starlike functions, Close to convex functions and bounded functions.

I. INTRODUCTION

Let \mathcal{A} denote the class of functions of the form $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ (1.1) which are analytic in the unit disc $\mathbb{E} = \{z: |z| < 1\}$. Let \mathcal{S} be the class of functions of the form (1.1), which are analytic univalent in \mathbb{E} .

In 1916, Bieber Bach ([2]) proved that $|a_2| \leq 2$ for the functions $f(z) \in \mathcal{S}$. In 1923, Löwner proved that $|a_3| \leq 3$ for the functions $f(z) \in \mathcal{S}$.

With the known estimates $|a_2| \leq 2$ and $|a_3| \leq 3$, it was natural to seek some relation between a_3 and a_2^2 for the class \mathcal{S} , Fekete and Szegö [5] used Löwner's method to prove the following well known result for the class \mathcal{S} .

Let $f(z) \in \mathcal{S}$, then

$$|a_3 - \mu a_2^2| \leq \begin{cases} 3 - 4\mu, & \text{if } \mu \leq 0; \\ 1 + 2 \exp\left(\frac{-2\mu}{1-\mu}\right), & \text{if } 0 \leq \mu \leq 1; \\ 4\mu - 3, & \text{if } \mu \geq 1. \end{cases} \quad (1.2)$$

The inequality (1.2) plays a very important role in determining estimates of higher coefficients for some sub classes \mathcal{S} (See Chhichra, Babalola). Several researchers proved many results for various classes and subclasses ([8]-[66]).

Let us define some subclasses of \mathcal{S} .

We denote by S^* , the class of univalent starlike functions

$$g(z) = z + \sum_{n=2}^{\infty} b_n z^n \in \mathcal{A} \text{ and satisfying the condition } \operatorname{Re} \left(\frac{zg'(z)}{g(z)} \right) > 0, z \in \mathbb{E}. \quad (1.3)$$

We denote by \mathcal{K} , the class of univalent convex functions $h(z) = z + \sum_{n=2}^{\infty} c_n z^n, z \in \mathcal{A}$ and satisfying the condition

$$\operatorname{Re} \left(\frac{zh'(z)}{h(z)} \right) > 0, z \in \mathbb{E}. \quad (1.4)$$

A function $f(z) \in \mathcal{A}$ is said to be close to convex if there exists $g(z) \in S^*$ such that

$$\operatorname{Re} \left(\frac{zf'(z)}{g(z)} \right) > 0, z \in \mathbb{E}. \quad (1.5)$$

The class of close to convex functions is denoted by \mathcal{C} and was introduced by Kaplan [3] and it was shown by him that all close to convex functions are univalent.

$$S^*(A, B) = \left\{ f(z) \in \mathcal{A}; \frac{zf'(z)}{f(z)} < \frac{1+Az}{1+Bz}, -1 \leq B < A \leq 1, z \in \mathbb{E} \right\} \quad (1.6)$$

$$\mathcal{K}(A, B) = \left\{ f(z) \in \mathcal{A}; \frac{(zf'(z))'}{f'(z)} < \frac{1+Az}{1+Bz}, -1 \leq B < A \leq 1, z \in \mathbb{E} \right\} \quad (1.7)$$

It is obvious that $S^*(A, B)$ is a subclass of S^* and $\mathcal{K}(A, B)$ is a subclass of \mathcal{K} .

We introduce a new subclass as

$$\left\{ f(z) \in \mathcal{A}; \alpha \frac{zf'(z)}{f(z)} + (1 - \alpha) \frac{zf(z)}{2 \int_0^z f(z) dz} < \frac{1+Az}{1+Bz}; z \in \mathbb{E} \right\}$$

and we will denote this class as $f(z) \in \mathcal{C}(S^*)^{-1}[A, B]$,

Symbol \prec stands for subordination, which we define as follows:

Principle of Subordination: Let $f(z)$ and $F(z)$ be two functions analytic in \mathbb{E} . Then $f(z)$ is called subordinate to $F(z)$ in \mathbb{E} if there exists a function $w(z)$ analytic in \mathbb{E} satisfying the conditions $w(0) = 0$ and $|w(z)| < 1$ such that $f(z) = F(w(z))$; $z \in \mathbb{E}$ and we write $f(z) \prec F(z)$. By \mathcal{U} , we denote the class of analytic bounded functions of the form $w(z) = \sum_{n=1}^{\infty} d_n z^n$, $w(0) = 0$, $|w(z)| < 1$. (1.8)
It is known that $|d_1| \leq 1$, $|d_2| \leq 1 - |d_1|^2$. (1.9)

II. PRELIMINARY LEMMAS

For $0 < c < 1$, we write $w(z) = \left(\frac{c+z}{1+cz}\right)$ so that $\frac{1+Aw(z)}{1+Bw(z)} = 1 + (A-B)c_1 z + (A-B)(c_2 - Bc_1^2)z^2 + \dots$ (2.1)

III. MAIN RESULT

THEOREM 3.1: Let $f(z) = z + \sum_{k=2}^{\infty} a_k z^k \in S^*(f, f', \alpha, \beta, \delta)$, then

$$|a_3 - \mu a_2^2| \leq \begin{cases} f[g - 4\mu\delta]\delta, & \text{if } \mu \leq w; \\ \frac{\delta}{3\alpha + \beta - 4\alpha\beta} & \text{if } w \leq \mu \leq y; \\ f[4\mu\delta - h]\delta, & \text{if } \mu \geq y \end{cases}$$

Where $f = \frac{1}{\{(1-\alpha)\beta + 2\alpha(1-\beta)\}^2}$
 $w = \frac{(8\alpha + 3\beta + 4\alpha^2 - 12\alpha^2\beta - 9\alpha\beta^2 - 3\alpha\beta + 9\alpha^2\beta^2)\delta - \{(1-\alpha)\beta + 2\alpha(1-\beta)\}^2}{4(3\alpha + \beta - 4\alpha\beta)\delta}$
 $y = \frac{(8\alpha + 3\beta + 4\alpha^2 - 12\alpha^2\beta - 9\alpha\beta^2 - 3\alpha\beta + 9\alpha^2\beta^2)\delta + \{(1-\alpha)\beta + 2\alpha(1-\beta)\}^2}{4(3\alpha + \beta - 4\alpha\beta)\delta}$
 $g = \frac{(8\alpha + 3\beta + 4\alpha^2 - 12\alpha^2\beta - 9\alpha\beta^2 - 3\alpha\beta + 9\alpha^2\beta^2)\delta - \{(1-\alpha)\beta + 2\alpha(1-\beta)\}^2}{(3\alpha + \beta - 4\alpha\beta)}$
 $h = \frac{(8\alpha + 3\beta + 4\alpha^2 - 12\alpha^2\beta - 9\alpha\beta^2 - 3\alpha\beta + 9\alpha^2\beta^2)\delta + \{(1-\alpha)\beta + 2\alpha(1-\beta)\}^2}{(3\alpha + \beta - 4\alpha\beta)}$

The result is sharp in the sense that right hand side is the least upper bound of the result and extremal function exists.

PROOF: By definition of $S^*(f, f', \alpha, \beta, \delta)$, we have

$$(1 - \alpha) \left(\frac{zf'(z)}{f(z)}\right)^\beta + \alpha \left(\frac{(zf'(z))^{1-\beta}}{f'(z)}\right) = \left(\frac{1+w(z)}{1-w(z)}\right)^\delta; w(z) \in \mathcal{U}. \tag{3.1}$$

Expanding the series (3.1), we get

$$(1 - \alpha) \left\{ 1 + \beta a_2 z + (2\beta a_3 + \frac{\beta(\beta-3)}{2} a_2^2) z^2 + \dots \right\} + \alpha \left\{ 1 + 2(1 - \beta) a_2 z + 2(1 - \beta)(3a_3 - (\beta + 2) a_2^2) z^2 + \dots \right\} = (1 + 2\delta c_1 z + 2\delta(c_2 + \delta c_1^2) z^2 + \dots). \tag{3.2}$$

Comparing the coefficients of like powers in equation (3.2), we get

$$a_2 = \frac{2\delta}{(1-\alpha)\beta + 2\alpha(1-\beta)} c_1 \text{ and } a_3 = \frac{\delta}{3\alpha + \beta - 4\alpha\beta} c_2 + \frac{8\alpha + 3\beta + 4\alpha^2 - 12\alpha^2\beta - 9\alpha\beta^2 - 3\alpha\beta + 9\alpha^2\beta^2}{(3\alpha + \beta - 4\alpha\beta)\{(1-\alpha)\beta + 2\alpha(1-\beta)\}^2} c_1^2 \delta^2. \tag{3.3}$$

From equation (3.3), we obtain

$$a_3 - \mu a_2^2 = \frac{\delta}{3\alpha + \beta - 4\alpha\beta} c_2 + \left[\frac{8\alpha + 3\beta + 4\alpha^2 - 12\alpha^2\beta - 9\alpha\beta^2 - 3\alpha\beta + 9\alpha^2\beta^2}{(3\alpha + \beta - 4\alpha\beta)\{(1-\alpha)\beta + 2\alpha(1-\beta)\}^2} - \frac{4}{\{(1-\alpha)\beta + 2\alpha(1-\beta)\}^2} \mu \right] \delta^2 c_1^2. \tag{3.4}$$

Taking absolute value, it can be rewritten as

$$|a_3 - \mu a_2^2| \leq \frac{\delta}{3\alpha + \beta - 4\alpha\beta} |c_2| + \frac{1}{\{(1-\alpha)\beta + 2\alpha(1-\beta)\}^2} \left| \frac{8\alpha + 3\beta + 4\alpha^2 - 12\alpha^2\beta - 9\alpha\beta^2 - 3\alpha\beta + 9\alpha^2\beta^2}{(3\alpha + \beta - 4\alpha\beta)} - 4\mu \right| \delta^2 |c_1^2|. \tag{3.5}$$

Using (1.9), we get

$$|a_3 - \mu a_2^2| \leq \frac{\delta}{3\alpha + \beta - 4\alpha\beta} (1 - |c_1|^2) + \frac{1}{\{(1-\alpha)\beta + 2\alpha(1-\beta)\}^2} \left| \frac{8\alpha + 3\beta + 4\alpha^2 - 12\alpha^2\beta - 9\alpha\beta^2 - 3\alpha\beta + 9\alpha^2\beta^2}{(3\alpha + \beta - 4\alpha\beta)} - 4\mu \right| |c_1^2| \delta^2 = \frac{\delta}{3\alpha + \beta - 4\alpha\beta} + \frac{1}{\{(1-\alpha)\beta + 2\alpha(1-\beta)\}^2} \left[\left| \frac{8\alpha + 3\beta + 4\alpha^2 - 12\alpha^2\beta - 9\alpha\beta^2 - 3\alpha\beta + 9\alpha^2\beta^2}{(3\alpha + \beta - 4\alpha\beta)} - 4\mu \right| \delta^2 - \frac{\{(1-\alpha)\beta + 2\alpha(1-\beta)\}^2 \delta}{3\alpha + \beta - 4\alpha\beta} \right] |c_1|^2.$$

Case I:

$$\mu \leq \frac{8\alpha+3\beta+4\alpha^2-12\alpha^2\beta-9\alpha\beta^2-3\alpha\beta+9\alpha^2\beta^2}{4(3\alpha+\beta-4\alpha\beta)}.$$

Above inequality can be rewritten as

$$\frac{|a_3 - \mu a_2^2|}{\delta} \leq \frac{1}{3\alpha+\beta-4\alpha\beta} + \frac{1}{\{(1-\alpha)\beta+2\alpha(1-\beta)\}^2} \left[\frac{(8\alpha+3\beta+4\alpha^2-12\alpha^2\beta-9\alpha\beta^2-3\alpha\beta+9\alpha^2\beta^2)\delta - \{(1-\alpha)\beta+2\alpha(1-\beta)\}^2}{(3\alpha+\beta-4\alpha\beta)} \right] \delta |c_1|^2 \quad (3.6)$$

Subcase I(a): $\mu \leq w$, where

$$w = \frac{(8\alpha+3\beta+4\alpha^2-12\alpha^2\beta-9\alpha\beta^2-3\alpha\beta+9\alpha^2\beta^2)\delta - \{(1-\alpha)\beta+2\alpha(1-\beta)\}^2}{4(3\alpha+\beta-4\alpha\beta)\delta}.$$

Using (1.9), we get from inequality (3.6) that

$$\frac{|a_3 - \mu a_2^2|}{\delta} \leq \frac{1}{\{(1-\alpha)\beta+2\alpha(1-\beta)\}^2} \left[\frac{(8\alpha+3\beta+4\alpha^2-12\alpha^2\beta-9\alpha\beta^2-3\alpha\beta+9\alpha^2\beta^2)\delta - \{(1-\alpha)\beta+2\alpha(1-\beta)\}^2}{(3\alpha+\beta-4\alpha\beta)} \right] \delta |c_1|^2 \quad (3.7)$$

Subcase I (b): $\mu \geq w$,

$$\text{where } w = \frac{(8\alpha+3\beta+4\alpha^2-12\alpha^2\beta-9\alpha\beta^2-3\alpha\beta+9\alpha^2\beta^2)\delta - \{(1-\alpha)\beta+2\alpha(1-\beta)\}^2}{4(3\alpha+\beta-4\alpha\beta)\delta}.$$

We obtain from inequality (3.6) that

$$|a_3 - \mu a_2^2| \leq \frac{\delta}{3\alpha+\beta-4\alpha\beta}. \quad (3.8)$$

Case II:

$$\mu \geq \frac{8\alpha + 3\beta + 4\alpha^2 - 12\alpha^2\beta - 9\alpha\beta^2 - 3\alpha\beta + 9\alpha^2\beta^2}{4(3\alpha + \beta - 4\alpha\beta)}$$

Proceeding as in case I, we get

$$\frac{|a_3 - \mu a_2^2|}{\delta} \leq \frac{\delta}{3\alpha+\beta-4\alpha\beta} + \frac{1}{\{(1-\alpha)\beta+2\alpha(1-\beta)\}^2} \left[4\mu\delta - \frac{(8\alpha+3\beta+4\alpha^2-12\alpha^2\beta-9\alpha\beta^2-3\alpha\beta+9\alpha^2\beta^2)\delta + \{(1-\alpha)\beta+2\alpha(1-\beta)\}^2}{(3\alpha+\beta-4\alpha\beta)} \right] \delta |c_1|^2 \quad (3.9)$$

Subcase II (a): $\mu \leq y$,

$$\text{where } y = \frac{(8\alpha+3\beta+4\alpha^2-12\alpha^2\beta-9\alpha\beta^2-3\alpha\beta+9\alpha^2\beta^2)\delta + \{(1-\alpha)\beta+2\alpha(1-\beta)\}^2}{4(3\alpha+\beta-4\alpha\beta)\delta}$$

It takes the form

$$|a_3 - \mu a_2^2| \leq \frac{\delta}{3\alpha+\beta-4\alpha\beta} \quad (3.10)$$

Combining results (3.8) of subcase I (b) and (3.10) of subcase II (a), we obtain

$$|a_3 - \mu a_2^2| \leq \frac{\delta}{3\alpha+\beta-4\alpha\beta} \quad \text{if } w \leq \mu \leq y \quad (3.11)$$

Where,

$$w = \frac{(8\alpha+3\beta+4\alpha^2-12\alpha^2\beta-9\alpha\beta^2-3\alpha\beta+9\alpha^2\beta^2)\delta - \{(1-\alpha)\beta+2\alpha(1-\beta)\}^2}{4(3\alpha+\beta-4\alpha\beta)\delta}$$

and

$$y = \frac{(8\alpha+3\beta+4\alpha^2-12\alpha^2\beta-9\alpha\beta^2-3\alpha\beta+9\alpha^2\beta^2)\delta + \{(1-\alpha)\beta+2\alpha(1-\beta)\}^2}{4(3\alpha+\beta-4\alpha\beta)\delta}$$

Subcase II (b): $\mu \geq y$,

$$\text{where, } y = \frac{(8\alpha+3\beta+4\alpha^2-12\alpha^2\beta-9\alpha\beta^2-3\alpha\beta+9\alpha^2\beta^2)\delta + \{(1-\alpha)\beta+2\alpha(1-\beta)\}^2}{4(3\alpha+\beta-4\alpha\beta)\delta}$$

Proceeding as in subcase I(a), we get

$$\frac{|a_3 - \mu a_2^2|}{\delta} \leq \frac{1}{\{(1-\alpha)\beta+2\alpha(1-\beta)\}^2} \left[4\mu\delta - \frac{(8\alpha+3\beta+4\alpha^2-12\alpha^2\beta-9\alpha\beta^2-3\alpha\beta+9\alpha^2\beta^2)\delta + \{(1-\alpha)\beta+2\alpha(1-\beta)\}^2}{(3\alpha+\beta-4\alpha\beta)} \right] \delta. \quad (3.12)$$

Hence the theorem is proved. The result is sharp in the sense that right hand side is the least upper bound of the result and extremal function exists. Extremal function for first and third inequality is defined by

$$f_1(z) = z \left\{ 1 + \sqrt{f}(2\delta - g)z \right\}^{\frac{2\delta}{2\delta-g}}$$

Extremal function for second inequality is defined by

$$f_2(z) = z \left(1 + \frac{1}{3\alpha+\beta-4\alpha\beta} z^2 \right)^\delta.$$

COROLLARY 3.2: Putting $\delta=1, \alpha = 1, \beta = 0$ in the theorem, we get

$$|a_3 - \mu a_2^2| \leq \begin{cases} 1 - \mu, & \text{if } \mu \leq \frac{2}{3}; \\ \frac{1}{3} \text{ if } \frac{2}{3} \leq \mu \leq \frac{4}{3}; \\ \mu - 1, & \text{if } \mu \geq \frac{4}{3} \end{cases}$$

These estimates were derived by Keogh and Merkes [8] and are results for the class of univalent convex functions.

COROLLARY 3.3: Putting $\delta=1, \alpha = 0, \beta = 1$ in the theorem, we get

$$|a_3 - \mu a_2^2| \leq \begin{cases} 3 - 4\mu, & \text{if } \mu \leq \frac{1}{2}; \\ \text{if } \frac{1}{2} \leq \mu \leq 1; \\ 4\mu - 3, & \text{if } \mu \geq 1 \end{cases}$$

These estimates were derived by Keogh and Merkes [8] and are results for the class of univalent starlike functions.

REFERENCES

- [1] Alexander, J.W, Function which map the interior of unit circle upon simple regions, Ann. Of Math., 17 (1995), 12-22.
- [2] Bieberbach, L., Uber die Koeffizienten derjenigen Potenzreihen, welche eine schlichte Abbildung des

- Einheitskrisisvermittlung, *S. – B. Preuss. Akad. Wiss.* 38 (1916), 940-955.
- [3] De Branges L., A proof of Bieberbach Conjecture, *Acta. Math.*, 154 (1985), 137-152.
- [4] Duren, P.L., Coefficient of univalent functions, *Bull. Amer. Math. Soc.*, 83 (1977), 891-911.
- [5] Fekete, M. and Szegő, G, Eine Bemerkung über ungeradeschlichtefunktionen, *J. London Math. Soc.*, 8 (1933), 85-89.
- [6] Garabedian, P. R., Schiffer, M., A Proof for the Bieberbach Conjecture for the fourth coefficient, *Arch. Rational Mech. Anal.*, 4 (1955), 427-465.
- [7] Kaur, C. and Singh, G., Approach to coefficient inequality for a new subclass of Starlike functions with extremals, *Int. Journal of Research in Advent Technology*, 5 (2017)
- [8] Kaur, C. and Singh, G., Coefficient Problem for A New Subclass of Analytic Functions Using Subordination, *International Journal of Research in Advent Technology*, 5 (2017)
- [9] Kaur, C. and Singh, G., Analytic Functions Subordinate To Leaf-Like Domain, *Advances in Mechanics*, 10 (1), 1444-1448, 2022
- [10] Kaur, G, Singh, G, Arif, M, Chinram, R, Iqbal, J, A study of third and fourth Hankel determinant problem for a particular class of bounded turning functions, *Mathematical Problems in Engineering*, 22, 511- 526, 2021
- [11] Keogh, F.R., Merkes, E.P., A coefficient inequality for certain classes of analytic functions, *Proc. Of Amer. Math. Soc.*, 20, 8-12, 1989.
- [12] Koebe, P., Über Die uniformisierungsbiliger analytischer Kurven, *Nach. Ges. Wiss. Gottingen* (1907), 633-669.
- [13] Lindelof, E., Memoire sur certaines inegalities dans la theorie des fonctions monogenes et sur quelques proprietes nouvelles de ces fonctions dans la voisinage d'un point singulier essentiel, *Acta Soc. Sci. Fenn.*, 23 (1909) , 481-519.
- [14] Ma, W. and Minda , D. Unified treatment of some special classes of univalent functions, Proceedings of the Conference on Complex Analysis , *Int. Press Tianjin* (1994) , 157-169.
- [15] Mehrotra, B. S, Singh, G, Saroa, M. S, Fekete-Szegő Inequality for Certain Subclasses of Analytic Functions, *Acta Ciencia Indica*, 39 (2), 97-104, 2013
- [16] Mehrotra, B. S, Singh, G, Saroa, M. S, Fekete-Szegő Inequality for a Certain Sub-classes of Analytic Functions, *Acta Ciencia Indica*, 39 (2), 125-138, 2013
- [17] Mehrotra, B. S, Singh, G, Saroa, M. S, Fekete-Szegő Inequality for a Certain Sub-classes of Analytic Functions, *Acta Ciencia Indica*, 39 (3), 217-228, 2013
- [18] Miller, S.S., Mocanu, P.T. And Reade, M.O., All convex functions are univalent and starlike, *Proc. of Amer. Math. Soc.*, 37 (1973), 553-554.
- [19] Nehari, Z. (1952), Conformal Mappings, McGraw- Hill, New York.
- [20] Nevanlinna, R., Über die Eigenschafteneiner analytischen Funktion in der Umgebung eines singulären steile order Linte, *Acta Soc. Sci. Fenn.*, 50 (1922), 1-46.
- [21] Pederson, R., A proof for the Bieberbach conjecture for the sixth coefficient, *Arch. Rational Mech. Anal.*, 31 (1968-69), 331-351.
- [22] Pederson, R. and Schiffer, M., A proof for the Bieberbach conjecture for the fifth coefficient, *Arch. Rational Mech. Anal.*, 45 (1972), 161-193.
- [23] Rani, M., Singh, G., Some classes of Schwarzian functions and its coefficient inequality that is sharp, *Turk. Jour. Of Comp. and Mathematics Education*, 11 (2020), 1366-1372.
- [24] Rathore, G. S., Singh, G. and Kumawat, L. et.al., Some Subclasses Of A New Class Of Analytic Functions under Fekete-Szegő Inequality, *Int. J. of Res. In Adv. Tech.*, 7 (2019)
- [25] Rathore, G. S., Singh, G., Fekete – Szegő Inequality for certain subclasses of analytic functions , *Journal Of Chemical , Biological And Physical Sciences*, 5 (2015) ,
- [26] Singh, G, Fekete – Szegő Inequality for a new class and its certain subclasses of analytic functions, *General Mathematical Notes*, 21 (2014),

- [27] Singh. G, Fekete – Szego Inequality for a new class of analytic functions and its subclass, *Mathematical Sciences International Research Journal*, 3 (2014),
- [28] Singh. G., Construction of Coefficient Inequality For a new Subclass of Class of Starlike Analytic Functions, *Russian Journal of Mathematical Research Series*, 1 (2015), 9-13.
- [29] Singh, G., Introduction of a new class of analytic functions with its Fekete–Szegő Inequality, *International Journal of Mathematical Archive*, 5 (2014), 30-35.
- [30] Singh. G, An inequality of second and third coefficients for a subclass of starlike functions constructed using nth derivative, *Kaav Int. J. of Sci. Eng. And Tech.*, 4 (2017), 206-210.
- [31] Singh, G, Fekete – Szego Inequality for asymptotic subclasses of family of analytic functions, *Stochastic Modelling and Applications*, 26 (2022),
- [32] Singh, G, Coefficient Inequality For Close To Starlike Functions Constructed Using Inverse Starlike Classes , *Kaav Int. J. of Sci. Eng. And Tech.*, 4 (2017), 177-182.
- [33] Singh. G, Coeff. Inequality for a subclass of Starlike functions that is constructed using nth derivative of the fns in the class, *Kaav Int. J. of Sci. Eng. And Tech.*, 4 (2017), 199-202.
- [34] Singh, G., Fekete – Szegő Inequality for functions approaching to a class in the limit form and another class directly, *Journal of Information and Computational Sciences*, 12 (4), 2022, 181-186
- [35] Singh, G., FeketeSzego Inequality For A Complicated Class Of Analytic Functions Approaching To A Class In The Limit Form And Other Class Directly, *Int. Journal of Research in Engineering and Science*, 10 (9), 619-624, 2022
- [36] Singh, G., Garg, J., Coefficient Inequality for A New Subclass of Analytic Functions, *Mathematical Sciences International Research Journal*, 4 (2015)
- [37] Singh, G, Singh, Gagan, Fekete – Szegő Inequality for Subclasses of A New Class of Analytic Functions, *Proceedings of the World Congress on Engineering*, (2014).
- [38] Singh, G, Saroa, M. S., and Mehrok, B. S., Fekete – Szegő Inequality For A New Class Of Analytic Functions , *Conference Of Information And Mathematical Sciences* , (2013).
- [39] Singh. G, Singh. Gagan, Saroa. M. S., Fekete – Szegő Inequality for a New Class of Convex Starlike Analytic Functions, *Conf. Of Information and Mathematical Sciences*, (2013).
- [40] Singh, G., Kaur, G., Coefficient Inequality for a Subclass of Starlike Function generated by symmetric points, *Ganita*, 70 (2020), 17-24.
- [41] Singh, G., Kaur, G., Coefficient Inequality for A New Subclass of Starlike Functions, *International Journal of Research in Advent Technology*, 5 (2017)
- [42] Singh, G., Kaur, G., Fekete-Szegő Inequality for A New Subclass of Starlike Functions, *International Journal of Research in Advent Technology*, 5 (2017)
- [43] Singh, G., Kaur, G., Fekete-Szegő Inequality for subclass of analytic function based on Generalized Derivative, *Aryabhata Journal Of Mathematics And Informatics*, 9 (2017)
- [44] Singh, G., Kaur, G., Coefficient Inequality For a subclass of analytic function using subordination method with extremal function, *Int. J. Of Adv. Res. in Sci&Engg*, 7 (2018)
- [45] Singh, G., Kaur, G., Fourth Hankel determinant for a new subclass of bounded turning functions, *Jnanabha*, 52 (1), 229-233, 2022
- [46] G Singh, G Singh, G Singh, A new subclass of univalent functions, *Уфимский математический журнал*, 11 (1), 132-139, 2019
- [47] Singh, G. and Kaur, G., 4th Hankel determinant for α bounded turning function, *Advances in Mathematics: Scientific Journal*, 9 (12), 10563-10567
- [48] Singh, G., Kaur, N., Fekete-Szegő Inequality for Certain Subclasses of Analytic Functions, *Mathematical Sciences International Research Journal*, 4 (2015)
- [49] Singh G, Saini G., Upper Bound Of Second Hankel Determinant Of Analytic Functions

- Constructed Using Composition *Stochastic Modelling & Applications*, 26 (3), 1411-1414, 2022
- [50] Singh, G, Singh, B, FeketeSzego coefficient inequality of regular functions for a special class, *Int. Journal of Research in Engineering and Science*, 10 (8), 2022, 556-560
- [51] Singh, G, Singh, B., FeketeSzego Coefficient Inequality Of Regular Functions For A Special Class, *Journal of Emerging Technologies and Innovative Research*, 9 (8), 625-632
- [52] Singh G., Konika, An interesting subclass constructed using a new parameter of an already defined class of regular functions and its coefficient inequality, *Int. Journal of Research in Engineering and Science*, 10 (9), 450-453
- [53] Singh G, Md AminulHoque, Fekete – Szego Inequality for asymptotic subclasses of family of analytic functions, *Stochastic Modelling And Applications*, 26 (8), 1464-1468, 2022
- [54] Singh G, Md AminulHoque, A specially designed class of univalent functions and establishment of its coefficient inequality, *NeuroQuantology* 20 (10), 2855-2859, 2022
- [55] Singh, G, Singh, P., Fekete-Szegő inequality for functions belonging to a certain class of analytic functions introduced using linear combination of variational powers of starlike and convex functions, *Journal Of Positive School Psychology*, 6 (2022), 8387-8391.
- [56] Singh, G, Singh, P., Fekete-Szegő Inequality For Class Of Close To Inverse Starlike Analytic Functions, *Advances in Mechanics* 10 (1), 1375-1386, 2022
- [57] Singh G., Patil A. S., An extraordinary class of asymptotic analytical functions with coefficient inequality, *NeuroQuantology*, 20(10), 4960-4966, 2022
- [58] Singh G., Bansal K., Kaur H., FeketeSzego Coefficient Inequality for a Subclass of Analytic Functions, *International Journal of Advances in Engineering and Management (IJAEM)*, 4 (10), 1295-1298, 2022
- [59] Singh. G, Rani M, An advance subclass of Analytic Functions having a unique coefficient inequality, *Int. J. of Res. in Engineering and Science*, 10 (8), 2022, 474-476
- [60] Singh. G, Rani M, On A New Problem Associated With P – Valent Starlike Functions, *Advances in Mechanics*, 10 (1), 1449-1456, 2022
- [61] Singh, G., Singh, G., Singh, G., A subclass of bi-univalent functions defined by generalized Sălăgean operator related to shell-like curves connected with Fibonacci numbers, *International Journal of Mathematics and Mathematical Sciences*, 2019
- [62] Singh G., Sharma N., Two new subclasses of already defined class of Analytic functions and establishment of their coefficient inequality, *NeuroQuantology*, 20(10), 4967-4976, 2022
- [63] Singh, G., Singh, G., Singh, G., Certain subclasses of univalent and bi-univalent functions related to shell-like curves connected with Fibonacci numbers, *General Mathematics*, 28 (1), 125-140, 2020
- [64] Singh, G., Singh, G., Singh, G., Certain subclasses of Sakaguchi type bi-univalent functions, *Ganita*, 69 (2), 45-55, 2019
- [65] Singh, G., Singh, G., Singh, G., Certain Subclasses of Bi-Close-to-Convex Functions Associated with Quasi Subordination, *Abstract and Applied Analysis*, 1, 1-6, 2019
- [66] Singh, G., Singh, G., Singh, G., Fourth Hankel determinant for a subclass of analytic functions defined by generalized Salagean operator, *Creat. Math. Inform.*, 31(2), 229-240, 2022
- [67] G Singh, G Singh, G Singh, Certain subclasses of multivalent functions defined with generalized Salagean operator and related to sigmoid function and lemniscate of Bernoulli, *J. Frac. Calc. Appl*, 13 (1), 65-81, 2022
- [68] G Singh, G Singh, G Singh, Upper bound on fourth Hankel determinant for certain subclass of multivalent functions, *Jnanabha*, 50 (2), 122-127, 2020
- [69] Srivastava H. M., G. Kaur, Singh. G, Estimates of fourth Hankel determinant for a class of analytic functions with bounded turnings involving cardioid domains, *Journal of*

Nonlinear and Convex Analysis, 22 (3), 511-526, 2022

- [70] SMC, P Hema, Gurmeet Singh, Coefficient Bounds Along with Their Extremal Functions Making Results Sharp For Certain Subclasses Of Analytic Functions Using Subordination Method, *Advances in Mechanics*, 10 (1), 1478-1486, 2022