Dental Anxiety and Fear: Practical Strategies to Help Patients Overcome Anxiety in Oral Healthcare

LARISSA BOM ROCCA LAPORT

Abstract- Dental anxiety and fear remain major barriers to accessing oral healthcare, often leading to delayed treatment, worsening of oral diseases, and reinforcement of avoidance behaviors. These conditions are shaped by previous negative experiences, anticipatory worry, and specific procedural triggers such as injections, drilling, or loss of control. Research consistently shows that effective management requires a multifaceted approach, integrating psychological, communicational, environmental, and pharmacological strategies. This article synthesizes current evidence on techniques for reducing dental anxiety, including the use of validated screening tools, empathic communication, patient-centered control, cognitive-behavioral interventions, environmental modifications, distraction methods, and evidence-based sedation practices. Emphasis is placed on tailoring interventions to individual triggers, implementing stepped-care models within dental practice, and fostering collaboration between dental teams and mental health professionals when necessary. By adopting systematic strategies to address anxiety, clinicians can reduce cancellations, improve adherence to treatment, and enhance long-term oral health outcomes.

Keywords: Dental Anxiety; Dental Fear; Patient Management; Cognitive-Behavioral Strategies; Sedation Dentistry; Patient-Centered Care; Oral Health Access.

I. INTRODUCTION

Dental anxiety is common across age groups and cultures and is consistently associated with avoidance of care, poorer oral health, and more emergency-driven treatment patterns. It is not a single construct but a cluster of anticipatory worry, situational fear (e.g., needles, drilling, gagging), and conditioned responses to earlier painful or humiliating encounters. A "vicious cycle" often develops: anxious patients postpone visits, oral diseases progress, treatment becomes more invasive, and fear is reinforced (Armfield, Stewart, & Spencer, 2007). Effective care therefore requires a dual aim: delivering technically sound dentistry while systematically reducing threat perception before, during, and after procedures.

The first step is to recognize and measure dental anxiety in routine practice. Brief, validated screening tools such as Corah's Dental Anxiety Scale and the Modified Dental Anxiety Scale (MDAS) can be completed in the waiting room and quickly flag patients who need tailored support (Corah, 1969; Humphris, Morrison, & Lindsay, 1995). Assessment should look beyond a total score to identify specific triggers—needles, loss of control, embarrassment about oral condition, or gag reflex—because targeted strategies are more effective than generic reassurance. For children and adolescents, screening also benefits from parental input, given that caregiver anxiety and modeling strongly shape young patients' responses to dental care (Gao et al., 2013).

Communication is the cornerstone of anxiety management. Empathic, nonjudgmental dialogue that validates fear ("many patients feel this way") reduces shame and opens space for collaborative planning. Clear explanation of procedures in plain language, calibrated to the individual's preference for detail, creates predictability. Techniques that enhance a patient's sense of control—such as pre-agreed stop signals, choice among acceptable options, and permission to pause—lower perceived threat and physiologic arousal. For children, the classic "Tell–Show–Do" approach remains foundational: describe what will happen, demonstrate with non-threatening models or on a finger, and then carry out the procedure, reinforcing coping behaviors throughout (Appukuttan, 2016). For adults with pronounced catastrophic thinking ("something terrible will happen," "I won't cope"), brief cognitive-behavioral elements—guided reappraisal of threat, coping self-statements, graduated exposure to feared stimuli, and post-visit debriefing—are strongly supported by evidence and can be integrated by trained dental teams or delivered via referral pathways (Wide Boman, Carlsson, Westin, & Hakeberg, 2013).

Environmental and sensory modifications further reduce arousal. Minimizing unpredictable noise, offering noise-cancelling headphones, and using neutral or pleasant scents help counter conditioned responses. Distraction strategies—music through headphones with patient-selected playlists, or immersive audiovisual content—reduce subjective anxiety and pain ratings during injections and operative care, with benefits documented in both adults and children (Lai et al., 2008; Appukuttan, 2016). Virtual reality and guided imagery show promise by absorbing attentional resources and situating the patient in soothing, controllable environments; these tools are particularly helpful during longer appointments that would otherwise deplete coping capacity (Appukuttan, 2016). For patients with strong reflexive responses, simple physiologic skills—paced breathing (e.g., 4–6 breaths per minute), grounding via tactile cues, and applied muscle tension for those prone to vasovagal syncope—are brief to teach and clinically useful.

Pain control is inseparable from anxiety control. Topical anesthetics with adequate onset time reduce needle-stick discomfort; warming and buffering local anesthetic, slow deposition, aspiration, and using smaller-gauge needles improve comfort and trust. Computer-assisted delivery systems can help certain patients who are highly sensitive to pressure changes. For those with a strong needle phobia, graded exposure away from the operatory—starting with viewing syringes, practicing coping statements and breathing, then rehearsal with a blunt tip on the arm before intraoral use—can convert avoidance into mastery over several short visits (Appukuttan, 2016). When indicated, nitrous oxide–oxygen inhalation sedation offers a rapid-onset, titratable, and quickly reversible option that alleviates anxiety while preserving protective reflexes and cooperation; careful screening and informed consent are essential. Short-acting oral benzodiazepines may be appropriate for procedural anxiolysis in selected adults, adhering strictly to sedation guidelines, dosing protocols, and escort requirements, and always combined with the behavioral measures noted above so patients develop durable coping skills rather than sole reliance on medication (Donaldson & Goodchild, 2015).

Clinical workflow can either amplify or ease anxiety. Scheduling anxious patients at times of day when the clinic is quieter and their personal stress is lower, starting with non-invasive or preventive services to "bank" positive experiences, and using shorter, more frequent appointments when stamina is limited are simple but high-yield adjustments. For restorative and surgical care, minimally invasive techniques—atraumatic restorative treatment, selective caries removal, rubber dam to control moisture and improve efficiency, and meticulous hemostasis—shorten chair time and reduce intraoperative stress. For children, pairing behavior guidance with topical fluoride, sealants, and silver diamine fluoride where appropriate can stabilize disease while trust and cooperative coping are built (Gao et al., 2013). In adults with long-standing avoidance and complex psychosocial histories, a staged care plan that begins with stabilization and desensitization before comprehensive rehabilitation often yields better adherence and outcomes.

Some patients present with severe dental phobia, comorbid psychiatric conditions, or trauma histories that make the operatory feel intrinsically unsafe. For these individuals, referral-based psychological care—most commonly cognitive-behavioral therapy (CBT) delivered by a trained therapist—has strong evidence for remission of dental phobia and improved dental attendance, with effects that endure beyond the end of therapy (Wide Boman et al., 2013). Hypnosis can be a useful adjunct for selected patients to modulate pain perception and anxiety during procedures, supported by meta-analytic evidence in medical settings, provided it is delivered by clinicians with appropriate training and within consented boundaries (Montgomery, Schnur, & David, 2011). A trauma-informed approach—anticipating triggers, avoiding coercive language, offering choices, and inviting consent at each stage—prevents re-traumatization and strengthens the therapeutic alliance. In every scenario, follow-up contact after challenging visits, brief reinforcement of coping successes, and transparent planning for the next appointment consolidate gains and reduce no-shows.

Implementation depends on team readiness. Brief training in recognizing dental anxiety, using MDAS screening, and delivering basic behavioral techniques can be incorporated into staff meetings and onboarding. Clinics can develop stepped-care pathways: (1) universal communication and sensory modifications for all; (2) targeted distraction, breathing skills, and enhanced local anesthesia for moderate anxiety; (3) nitrous oxide or minimal oral sedation plus structured CBT elements for high anxiety; and (4) referral for specialized CBT or, when necessary, advanced sedation within appropriately equipped settings. Recording anxiety scores, triggers, and successful strategies in the chart allows continuity across providers and supports quality improvement. Over time, such systems reduce cancellations, shorten procedure times, improve patient satisfaction, and foster equity by making care accessible to individuals who have historically avoided dentistry due to fear (Armfield et al., 2007; Appukuttan, 2016).

The flowchart illustrates a stepwise approach to overcoming dental anxiety and fear, beginning with initial screening to identify anxious patients early. It then emphasizes empathetic communication, ensuring patients feel understood and supported, followed by patient-centered control, where individuals are given choices and a sense of autonomy during treatment. Next, environmental and distraction techniques such as calming surroundings and sensory modifications are applied to reduce stress. Finally, when necessary, pharmacological interventions like sedation or anxiolytics are introduced as complementary strategies. Together, these stages highlight a comprehensive, patient-focused framework for managing dental anxiety effectively.



Figure 1. Flowchart of Strategies to Overcome Dental Anxiety and Fear

Source: Created by author.

In sum, dental anxiety is both prevalent and modifiable. When clinicians pair empathic communication and evidence-based behavioral techniques with excellent analgesia and judicious use of sedation, patients learn to anticipate—not dread—dental visits. Measured routinely, addressed proactively, and followed up deliberately, anxiety becomes another clinical parameter that teams skillfully manage, closing the loop on the vicious cycle and opening a path to sustained oral health.

REFERENCES

[1] Appukuttan, D. (2016). Strategies to manage patients with dental anxiety and dental phobia: Literature review. Clinical, Cosmetic and Investigational Dentistry, 8, 35–50.

[2] Armfield, J. M., Stewart, J. F., & Spencer, A. J. (2007). The vicious cycle of dental fear: Exploring the interplay between oral health, service utilization and dental fear. BMC Oral Health, 7, 1.

[3] Corah, N. L. (1969). Development of a dental anxiety scale. Journal of Dental Research, 48, 596.

[4] Donaldson, M., & Goodchild, J. H. (2015). Appropriate oral sedation in the dental office. Dental Clinics of North America, 59(2), 389–406.

[5] Gao, X., Hamzah, S. H., Yiu, C. K. Y., McGrath, C., & King, N. M. (2013). Dental fear and anxiety in children and adolescents: Qualitative studies using content analysis. Journal of Dental Research, 92(7), 600–606.

[6] Humphris, G. M., Morrison, T., & Lindsay, S. J. (1995). The Modified Dental Anxiety Scale: UK norms and evidence for validity. Community Dental Health, 12(3), 143–150.

[7] Lai, H. L., Chen, C. J., Peng, T. C., Chang, F. M., Hsieh, Y. M., & Huang, H. Y. (2008). Randomized controlled trial of music during dental treatments to reduce anxiety. Journal of Clinical Nursing, 17(19), 2654–2660.

[8] Montgomery, G. H., Schnur, J. B., & David, D. (2011). The impact of hypnotic suggestion on clinical pain: Meta-analysis. International Journal of Clinical and Experimental Hypnosis, 59(3), 339–357.

[9] Wide Boman, U., Carlsson, V., Westin, M., & Hakeberg, M. (2013). Psychological treatment of dental anxiety among adults: A systematic review. European Journal of Oral Sciences, 121(3), 225–234.

[10] SANTOS,Hugo;PESSOA,EliomarGotardi.Impactsofdigitalizationontheefficiencyandqualityofpublicservices:Acomprehensiveanalysis.LUMENETVIRTUS,[S.l.],v.15,n.40,p.44094414,2024.DOI:10.56238/levv15n40024.Disponívelem:https://periodicos.newsciencepubl.com/LEV/article/view/452.Acessoem:25jan.2025.

[11] Freitas,G.B.,Rabelo,E.M.,&Pessoa,E.G.(2023). Projetomodularcomreaproveitamentodecontainermaritimo.BrazilianJournalofDevelopment,9(10),28303–28339.https://doi.org/10.34117/bjdv9n10057

[12] Freitas,G.B.,Rabelo,E.M.,&Pessoa,E.G.(2023). Projetomodularcomreaproveitamentodecontainermaritimo.BrazilianJournalofDevelopment,9(10),28303–28339.https://doi.org/10.34117/bjdv9n10057

[13] Pessoa,E.G.,Feitosa,L.M.,ePadua,V.P.,&Pereira,A.G.(2023).Estudodosrecalquesprimáriosemumaterroexecutadosobreaargilamoledo Sarapuí.BrazilianJournalofDevelopment,9(10),28352–28375.https://doi.org/10.34117/bjdv9n10059

[14] PESSOA,E.G.;FEITOSA,L.M.;PEREIRA,A.G.;EPADUA,V.P.Efeitosdeespéciesdealnaeficiênciadecoagulação,Alresidualepropriedadedosflocosnotratamentodeáguassuperficiais.BrazilianJournalofHealthReview,[S.l.],v.6,n.5,p.2481424826,2023.DOI:10.34119/bjhrv6n5523.Disponívelem:https://ojs.brazilianjournals.com.br/ojs/index.php/BJHR/article/view/63890.Acessoem:25jan.2025.

[15] SANTOS,Hugo;PESSOA,EliomarGotardi.Impactsofdigitalizationontheefficiencyandqualityofpublicservices:Acomprehensiveanalysis.LUMENETVIRTUS,[S.l.],v.15,n.40,p.44094414,2024.DOI:10.56238/levv15n40024.Disponívelem:https://periodicos.newsciencepubl.com/LEV/article/view/452.Acessoem:25jan.2025.

[16] Filho, W. L. R. (2025). The Role of Zero Trust Architecture in Modern Cybersecurity: Integration with IAM and Emerging Technologies. *Brazilian Journal of Development*, *11*(1), e76836. https://doi.org/10.34117/bjdv11n1-060

[17] Oliveira, C. E. C. de. (2025). Gentrification, urban revitalization, and social equity: challenges and solutions. *Brazilian Journal of Development*, *11*(2), e77293. https://doi.org/10.34117/bjdv11n2-010

[18] Pessoa, E. G. (2024). Pavimentos permeáveis uma solução sustentável. *Revista Sistemática*, *14*(3), 594–599. https://doi.org/10.56238/rcsv14n3-012

[19] Filho, W. L. R. (2025). THE ROLE OF AI IN ENHANCING IDENTITY AND ACCESS MANAGEMENT SYSTEMS. *International Seven Journal of Multidisciplinary*, *1*(2). https://doi.org/10.56238/isevmjv1n2-011

[20] Antonio, S. L. (2025). Technological innovations and geomechanical challenges in Midland Basin Drilling. Brazilian Journal of Development, 11(3), e78097. https://doi.org/10.34117/bjdv11n3-005

[21] Pessoa, E. G. (2024). Pavimentos permeáveis uma solução sustentável. *Revista Sistemática*, *14*(3), 594–599. https://doi.org/10.56238/rcsv14n3-012

[22] Pessoa, E. G. (2024). Pavimentos permeáveis uma solução sustentável. *Revista Sistemática*, *14*(3), 594–599. https://doi.org/10.56238/rcsv14n3-012
[23] Eliomar Gotardi Pessoa, & Coautora: Glaucia Brandão Freitas. (2022). ANÁLISE DE CUSTO DE PAVIMENTOS PERMEÁVEIS EM BLOCO DE CONCRETO UTILIZANDO BIM (BUILDING INFORMATION MODELING). Revistaft, 26(111), 86. https://doi.org/10.5281/zenodo.10022486
[24] Eliomar Gotardi Pessoa, Gabriel Seixas Pinto Azevedo Benittez, Nathalia Pizzol de Oliveira, & Vitor Borges Ferreira Leite. (2022). ANÁLISE COMPARATIVA ENTRE RESULTADOS EXPERIMENTAIS E TEÓRICOS DE UMA ESTACA COM CARGA HORIZONTAL APLICADA NO TOPO. Revistaft, 27(119), 67. https://doi.org/10.5281/zenodo.7626667
[25] Eliomar Gotardi Pessoa, & Coautora: Glaucia Brandão Freitas. (2022). ANÁLISE COMPARATIVA ENTRE RESULTADOS TEÓRICOS DA DEFLEXÃO DE UMA LAJE PLANA COM CARGA DISTRIBUÍDA PELO MÉTODO DE EQUAÇÃO DE DIFERENCIAL DE LAGRANGE POR SÉRIE DE FOURIER DUPLA E MODELAGEM NUMÉRICA PELO SOFTWARE SAP2000. Revistaft, 26(111), 43. https://doi.org/10.5281/zenodo.10019943
[26] Pessoa, E. G. (2025). Optimizing helical pile foundations: a comprehensive study on displaced soil volume and group behavior. *Brazilian Journal of Development*, *11*(4), e79278. https://doi.org/10.34117/bjdv11n4-047
[27] Pessoa, E. G. (2025). Utilizing recycled construction and demolition waste in permeable pavements for sustainable urban infrastructure. *Brazilian Journal of Development*, *11*(4), e79277. https://doi.org/10.34117/bjdv11n4-046