

Herbal Formulation for Hair Growth Promotion and Scalp Health

SHINDE TAJAS SANJAY¹, BHUSARE SUSHANT BHASKAR², TUPE ROHIT ROHIDAS³, GADE DHANANJAY ASMAN⁴

^{1, 2, 3, 4}Department of Pharmacognosy / Pharmaceutical Sciences, HSBPVT'S Faculty of Pharmacy, Kashti., Kashti, Shrigonda, India

Abstract- Polyherbal hair oils represent one of the most widely used categories in herbal cosmetic science, combining multiple medicinal plant extracts to achieve synergistic therapeutic effects for hair growth promotion and scalp health improvement. This review comprehensively examines the botanical constituents, phytochemical profiles, mechanisms of action, formulation methods, and evaluation parameters relevant to polyherbal hair oil preparations. Key medicinal plants—Amla (*Embllica officinalis*), Bhringraj (*Eclipta alba*), Neem (*Azadirachta indica*), Hibiscus (*Hibiscus rosa-sinensis*), and Fenugreek (*Trigonella foenum-graecum*)—are discussed in detail with respect to their documented pharmacological activities including antioxidant, antimicrobial, anti-inflammatory, and hair growth-stimulating properties. The review further addresses common hair disorders such as alopecia, dandruff, premature greying, and hair thinning, and discusses how polyherbal formulations target these conditions through multi-modal biological pathways. Formulation methodology, standardized quality evaluation parameters, stability testing, and recent advances in nano-herbal delivery systems are systematically reviewed. The study concludes that polyherbal hair oils offer a scientifically validated, economically accessible, and environmentally sustainable alternative to synthetic cosmetic products, with significant potential for industrial-scale standardization and clinical application.

Keywords: Polyherbal hair oil, Amla, Bhringraj, Neem, Hibiscus, Fenugreek, Hair growth, Alopecia, Phytoconstituents, Herbal cosmetics, Scalp health, Antidandruff

I. INTRODUCTION

Hair is a keratinous filamentous structure that plays a significant role in the aesthetic appearance, social identity, and psychological well-being of individuals across all cultures. Beyond its cosmetic relevance, hair serves physiological functions including

thermoregulation, sensory perception, and UV radiation protection. In contemporary society, hair disorders such as alopecia, excessive hair fall, dandruff, and premature greying have become increasingly prevalent owing to multifactorial causes including environmental pollution, chronic stress, nutritional deficiencies, hormonal dysregulation, and the indiscriminate use of chemically aggressive hair care products.

The global herbal cosmetics market has witnessed exponential growth driven by increasing consumer awareness of the adverse effects associated with synthetic chemical formulations. Herbal hair oils, derived from medicinal plants with documented therapeutic properties, offer a safer, multifunctional, and holistic approach to hair and scalp care. The use of plant-based oils and botanical extracts in hair care is deeply embedded in traditional medicine systems including Ayurveda, Unani, and Traditional Chinese Medicine, where preparations combining multiple herbs are preferred over single-herb formulations due to their synergistic bioactivity.

Polyherbal formulations—those incorporating two or more botanical ingredients—leverage the combined phytochemical profiles of individual plants to produce additive or synergistic pharmacological effects. This principle, known as polypharmacy in traditional medicine, has been validated by numerous in vitro and in vivo studies demonstrating enhanced efficacy compared to single-herb extracts. This review systematically evaluates the scientific evidence supporting polyherbal hair oil formulations, covering botanical characterization, phytochemistry, mechanisms of therapeutic action, formulation science, quality evaluation, and future research directions.

II. HAIR ANATOMY AND PHYSIOLOGY

A thorough understanding of hair structure and the hair growth cycle is foundational to appreciating the mechanisms by which polyherbal formulations exert their therapeutic effects. Hair is composed of two structurally distinct components: the hair shaft (the visible portion above the skin surface) and the hair follicle (the dynamic structure embedded in the dermis).

2.1 Structural Organization of Hair

The hair shaft consists of three concentric layers. The medulla, the innermost core, is present in thick hair types and is composed of loosely organized cells. The cortex, constituting the bulk of the hair shaft, comprises densely packed keratin filaments and melanin granules that determine hair color, strength, and elasticity. The cuticle, the outermost protective layer, is formed by overlapping scale-like cells and is responsible for hair's sheen, smoothness, and moisture retention capacity.

The hair follicle is a complex mini-organ containing the hair bulb at its base, which houses the matrix cells responsible for hair growth, the dermal papilla (a highly vascular, innervated connective tissue structure that regulates follicular activity), and the outer and inner root sheaths. Associated with the follicle are the arrector pili muscle and sebaceous glands, which contribute to scalp lubrication and moisture balance.

2.2 Hair Growth Cycle

Hair growth proceeds through three distinct and cyclically recurring phases:

- **Anagen (Active Growth Phase):** The longest phase, lasting 2–7 years, during which matrix cells in the hair bulb actively proliferate and differentiate to produce the hair shaft. Approximately 85–90% of scalp hairs are in the anagen phase at any given time.
- **Catagen (Transitional Phase):** A brief phase of 2–4 weeks characterized by cessation of hair growth, apoptosis of matrix cells, regression of the hair bulb, and detachment of the follicle from the dermal papilla.
- **Telogen (Resting Phase):** Lasting 3–5 months, this quiescent phase culminates in the shedding of

the club hair and the initiation of a new anagen phase. Disruption of anagen-to-telogen ratio, as seen in telogen effluvium, directly contributes to increased hair shedding.

Disorders affecting hair growth typically involve premature entry into the catagen phase, prolonged telogen duration, miniaturization of follicles, or destruction of follicular stem cells. Polyherbal hair oils target these pathways through multiple phytochemical mechanisms.

III. COMMON HAIR DISORDERS

Hair disorders represent a diverse group of conditions affecting the hair shaft, follicle, and scalp. Their etiology is multifactorial, involving genetic predisposition, endocrine dysregulation, nutritional imbalances, infectious agents, and environmental factors.

Androgenetic alopecia (AGA), the most common form of hair loss affecting both males and females, is mediated by the action of dihydrotestosterone (DHT) on genetically susceptible follicles, leading to progressive follicular miniaturization and eventual hair loss. Telogen effluvium, typically triggered by physiological or psychological stress, is characterized by a diffuse, non-scarring hair loss due to premature follicular entry into the telogen phase. Alopecia areata, an autoimmune condition, involves T-lymphocyte-mediated destruction of hair follicles, resulting in patchy hair loss.

Dandruff (seborrheic dermatitis), characterized by excessive desquamation of scalp corneocytes and associated with *Malassezia furfur* colonization, affects up to 50% of the global adult population. Premature canities (greying) results from melanocyte depletion or dysfunction in the hair bulb due to oxidative stress, genetic factors, or nutritional deficiencies. Additional conditions including trichorrhexis nodosa, split ends, and scalp xerosis are commonly exacerbated by chemical processing, heat styling, and UV exposure.

IV. REVIEW OF KEY HERBAL INGREDIENTS

formulations, their pharmacognostic sources, major phytoconstituents, and documented therapeutic activities:

The following table summarizes the principal botanical ingredients used in polyherbal hair oil

Table 1: Key Herbal Ingredients, Phytoconstituents, and Therapeutic Activities

Herbal Ingredient	Part Used	Key Phytoconstituents	Therapeutic Activity
Amla (<i>Emblica officinalis</i>)	Fruit	Vitamin C, Tannins, Flavonoids, Gallic acid	Antioxidant, hair strengthening, anti-greying
Bhringraj (<i>Eclipta alba</i>)	Whole plant	Wedelolactone, Ecliptine, Flavonoids	Hair growth stimulation, anti-alopecia, anti-inflammatory
Neem (<i>Azadirachta indica</i>)	Leaves, Oil	Nimbidin, Azadirachtin, Nimbin, Quercetin	Antimicrobial, antifungal, antidandruff
Hibiscus (<i>Hibiscus rosa-sinensis</i>)	Flowers, Leaves	Anthocyanins, Mucilage, Flavonoids	Hair conditioning, softness, reduces hair fall
Fenugreek (<i>Trigonella foenum-graecum</i>)	Seeds	Trigonelline, Diosgenin, Mucilage, Saponins	Nourishes follicles, reduces breakage, protein-rich
Coconut Oil (<i>Cocos nucifera</i>)	Fruit (oil)	Lauric acid, Caprylic acid, Medium-chain fatty acids	Carrier oil, deep penetration, moisture retention
Brahmi (<i>Bacopa monnieri</i>)	Leaves	Bacosides A & B, Alkaloids, Saponins	Stress reduction, scalp nourishment, cooling effect

4.1 Amla (*Emblica officinalis* Gaertn.)

Amla, the Indian gooseberry, is one of the most extensively studied plants in Ayurvedic medicine. Its fruits contain exceptionally high concentrations of ascorbic acid (Vitamin C) — one of the richest natural sources known — along with tannins (emblicanin A and B), gallic acid, ellagic acid, and flavonoids. These constituents confer potent antioxidant activity, scavenging free radicals that contribute to oxidative damage of hair follicles and premature greying. Amla extracts have been demonstrated to inhibit 5-alpha reductase activity, thereby reducing DHT-mediated follicular miniaturization. Additionally, the tannin content imparts astringent properties that strengthen hair roots and reduce hair breakage.

4.2 Bhringraj (*Eclipta alba* Hassk.)

Bhringraj, regarded in Ayurveda as the 'king of herbs' for hair care (Kesharaja), has been extensively

validated for its hair growth-stimulating properties. The plant contains wedelolactone, ecliptine, luteolin, and coumestans as principal bioactive constituents. Pharmacological studies have demonstrated that methanolic extracts of *Eclipta alba* significantly accelerate anagen phase initiation and prolong anagen duration in murine hair growth models, with efficacy comparable to minoxidil 2%. The mechanisms involve upregulation of Wnt/beta-catenin signaling in follicular keratinocytes and increased dermal papilla proliferation. Anti-inflammatory and immunomodulatory effects further contribute to its therapeutic profile.

4.3 Neem (*Azadirachta indica* A. Juss.)

Neem has been documented in traditional systems of medicine for over 4,000 years and is recognized globally for its broad-spectrum antimicrobial properties. The principal bioactive limonoids—azadirachtin, nimbidin, nimbin, and nimbolide—

exhibit potent antifungal activity against *Malassezia furfur*, the primary etiological agent of dandruff and seborrheic dermatitis, as well as antibacterial activity against *Staphylococcus aureus* and *Pseudomonas aeruginosa* implicated in scalp folliculitis. Neem oil also contains fatty acids (oleic, linoleic, stearic) that contribute to scalp moisturization and barrier function restoration.

4.4 Hibiscus (*Hibiscus rosa-sinensis* L.)

Both flowers and leaves of *Hibiscus rosa-sinensis* are utilized in traditional hair care preparations. The flowers are rich in anthocyanins, flavonoids (quercetin, kaempferol), and mucilage polysaccharides, while the leaves contain protein-rich mucilaginous compounds. Experimental studies indicate that hibiscus extracts promote hair follicle proliferation and increase anagen-phase follicle density. The conditioning properties of hibiscus, mediated by its mucilage content and film-forming capacity, improve hair manageability, reduce combing friction, and enhance shine. Anti-allopecic activity has been attributed to selective inhibition of 5- α reductase.

4.5 Fenugreek (*Trigonella foenum-graecum* L.)

Fenugreek seeds contain a unique combination of steroidal saponins (diosgenin, tigogenin), trigonelline (an alkaloid precursor to nicotinic acid/Vitamin B3), galactomannan mucilage, and proteins rich in lysine and tryptophan. The high protein content directly nourishes hair keratin structure, reducing breakage and improving tensile strength. Diosgenin exhibits weak estrogenic activity and has been proposed as a modulator of androgenic alopecia pathways. The lecithin content in fenugreek seeds provides deep conditioning and is implicated in improved moisture retention within the hair shaft.

V. PHYTOCHEMICAL CONSTITUENTS AND MECHANISMS OF ACTION

5.1 Key Phytochemical Classes

The therapeutic activity of polyherbal hair oils is attributed to complex interactions among diverse classes of secondary metabolites present in the constituent plant materials:

- **Flavonoids and Polyphenols:** Quercetin, kaempferol, rutin, and gallic acid provide

antioxidant protection to follicular cells, scavenging reactive oxygen species (ROS) that damage DNA and disrupt normal cell division in the hair matrix.

- **Tannins:** Hydrolyzable tannins (gallotannins, ellagitannins) from Amla and other sources confer astringent properties, tighten scalp pores, and strengthen hair follicular attachment to the scalp dermis.
- **Alkaloids:** Trigonelline (fenugreek) and ecliptine (bhringraj) modulate cellular signaling pathways involved in follicular cycling and proliferation.
- **Saponins:** Exhibit surfactant and cleansing properties, remove excess sebum without disrupting the scalp's natural lipid barrier.
- **Mucilage Polysaccharides:** Form a protective film over the hair shaft, improve moisture retention, reduce friction and static charge, and enhance conditioning and manageability.
- **Anthocyanins:** Pigmented flavonoids with exceptional radical-scavenging capacity, contributing to anti-greying and cytoprotective effects.
- **Fatty Acids:** Saturated (lauric acid in coconut oil) and unsaturated (oleic, linoleic) fatty acids penetrate the hair cortex to strengthen the internal structure and reduce protein loss.

5.2 Mechanisms of Therapeutic Action

Polyherbal hair oils exert their beneficial effects through several synergistic biological mechanisms:

1. **Enhanced Scalp Microcirculation:** Vasodilatory phytoconstituents (wedelolactone, flavonoids) promote blood flow to the dermal papilla, improving nutrient and oxygen delivery to proliferating follicular matrix cells.
2. **Follicular Stem Cell Activation:** Wnt/ β -catenin signaling activation by *Eclipta alba* extracts stimulates follicular stem cells in the bulge region, facilitating anagen re-entry after telogen.
3. **5- α Reductase Inhibition:** Polyphenols and steroidal saponins from Amla, Fenugreek, and Hibiscus competitively inhibit the enzyme responsible for converting testosterone to the more potent hair-follicle-miniaturizing DHT.
4. **Anti-inflammatory Activity:** Nimbidin (neem), wedelolactone (bhringraj), and flavonoids broadly inhibit pro-inflammatory cytokines (IL-1 β , TNF-

- α) that compromise follicular health and contribute to inflammatory alopecia.
5. Antimicrobial and Antifungal Effects: Azadirachtin and neem limonoids disrupt fungal cell membranes and inhibit sterol biosynthesis in *Malassezia* spp., addressing the root cause of dandruff.
 6. Antioxidant Cytoprotection: High-potency antioxidants (Vitamin C from Amla, anthocyanins from Hibiscus) neutralize oxidative stress in melanocytes and follicular cells, slowing premature greying and follicular aging.
 7. Moisture Retention and Conditioning: Mucilaginous compounds and fatty acids form moisture-retaining films around the hair shaft, preventing hygral fatigue and maintaining optimal hair fiber hydration.

VI. FORMULATION METHODOLOGY

The preparation of polyherbal hair oil is a multi-step process requiring careful selection of raw materials, standardized extraction procedures, appropriate oil base selection, and controlled processing conditions to maximize bioactive retention while ensuring product stability.

6.1 Ingredient Selection and Preparation

Herbal raw materials are selected based on pharmacognostic identity, purity, and freedom from pesticide contamination. After authentication (macroscopic, microscopic, and chemotaxonomic verification), plant materials are shade-dried at temperatures not exceeding 40°C to preserve thermolabile phytoconstituents such as Vitamin C and volatile terpenes. Dried materials are coarsely powdered and stored in airtight, moisture-proof containers until use.

6.2 Oil Base Selection

The carrier oil is a critical component of the formulation, serving both as an extraction medium and as a therapeutic ingredient. Coconut oil (*Cocos nucifera*) is most frequently employed due to its high lauric acid content, which penetrates the hair cortex and reduces protein loss. Sesame oil (*Sesamum indicum*) provides UV-absorbing lignans (sesamol, sesamin) and excellent skin permeation. Castor oil (*Ricinus communis*), with its high ricinoleic acid

content, enhances hair fiber flexibility and shine. Base oil selection influences viscosity, spreading coefficient, and bioactive extractability.

6.3 Hot Infusion (Murchhana) Method

The hot infusion or murchhana method, derived from Ayurvedic pharmaceutical tradition, is the most widely employed preparation technique. Accurately weighed powdered herbal materials are combined with measured volumes of oil base in a suitable vessel. The mixture is heated to 60–80°C under continuous stirring for a specified duration (typically 4–8 hours), allowing phytoconstituent extraction from the plant matrix into the lipophilic oil phase. Temperature control is critical — excessive heat degrades thermosensitive bioactives, while insufficient temperatures result in incomplete extraction.

6.4 Filtration and Finishing

Following the extraction phase, the oil is cooled to room temperature and filtered through multiple layers of muslin cloth or a Buckner funnel with filter paper to remove all particulate plant material. Permitted preservatives (tocopherol at 0.1–0.5%, BHT at <0.01%) may be incorporated at this stage to extend shelf life. The finished product is filled into amber-colored glass or UV-resistant plastic bottles under nitrogen headspace to minimize oxidative degradation, labeled with batch number and expiry date, and stored at ambient conditions (15–30°C) away from direct light and moisture.

VII. EVALUATION PARAMETERS

Rigorous physicochemical, microbiological, and stability evaluation is essential for establishing quality standards, batch-to-batch consistency, and therapeutic reproducibility of polyherbal hair oil formulations. The following table summarizes the principal evaluation parameters and their significance:

Table 2: Evaluation Parameters, Methods, and Acceptable Standards for Polyherbal Hair Oil

Evaluation Parameter	Method/Test	Acceptable Range / Significance
Organoleptic Properties	Visual inspection, olfactometry	Color, odor, clarity, texture
pH	Digital pH meter (10% w/v emulsion)	4.5 – 6.5 (slightly acidic for scalp compatibility)
Viscosity	Brookfield viscometer at 25°C	Consistent flow; should match formulation standard
Acid Value	Titration with KOH	Low acid value indicates oil stability
Specific Gravity	Specific gravity bottle at 25°C	Consistent density; usually 0.85–0.95 g/mL
Antimicrobial Activity	Agar well diffusion / disc diffusion	Zone of inhibition against <i>S. aureus</i> , <i>C. albicans</i>
Stability Testing	ICH guidelines (40°C/75% RH, 6 months)	No phase separation, rancidity, or color change
Spreadability	Parallel plate method	Good spreadability indicates ease of application

The pH of the formulation is particularly critical, as the scalp's natural pH of 4.5–5.5 is essential for maintaining the integrity of the cuticle layer and deterring microbial colonization. Formulations deviating significantly from this range may cause cuticle lifting, increased friction, and protein denaturation. Stability studies conducted per ICH

Q1A guidelines (40°C/75% relative humidity for 6 months) assess accelerated degradation and determine appropriate shelf life. Organoleptic consistency — uniform color, characteristic herbal odor, and appropriate viscosity — serves as a primary quality control indicator.

VIII. THERAPEUTIC APPLICATIONS

Polyherbal hair oils demonstrate a broad spectrum of therapeutic applications that address both the cosmetic and clinical dimensions of hair and scalp health:

- **Hair Growth Promotion:** Through 5-alpha reductase inhibition, Wnt pathway activation, and improved dermal papilla vascularization, polyherbal oils effectively stimulate new hair growth and prevent follicular miniaturization in androgenetic alopecia and stress-induced hair loss.
- **Anti-Dandruff Activity:** The antifungal properties of neem constituents against *Malassezia furfur*, combined with anti-inflammatory effects, address both the infectious and inflammatory components of seborrheic dermatitis.
- **Reduction of Hair Fall:** Strengthening of the hair follicular attachment zone, improvement in cortical protein structure, and reduction of oxidative follicular damage collectively reduce daily hair shedding.
- **Scalp Health Improvement:** Antimicrobial, anti-inflammatory, and sebum-balancing properties normalize scalp microbiome composition and barrier function.
- **Enhancement of Hair Texture and Shine:** Conditioning mucilage, proteins, and fatty acids smooth the cuticle layer, reflecting light uniformly to improve hair sheen and reduce frizz.
- **Prevention of Premature Greying:** Antioxidant protection of follicular melanocytes by polyphenols and Vitamin C slows the age-related and stress-induced loss of hair pigmentation.

IX. ADVANTAGES OVER SYNTHETIC FORMULATIONS

The growing preference for herbal hair care products is underpinned by several documented advantages over conventional synthetic formulations:

- **Safety and Tolerability:** Herbal ingredients have millennia of documented human use with well-characterized safety profiles. Unlike synthetic agents such as minoxidil, which may cause scalp irritation, contact dermatitis, and systemic side effects, polyherbal formulations generally demonstrate excellent tolerability.
- **Multifunctional Synergy:** The complex phytochemical matrix of each herbal ingredient, and the combined phytochemistry across multiple plants, provides simultaneous action on multiple pathogenic pathways—an advantage no single synthetic molecule can replicate.
- **Biodegradability and Environmental Sustainability:** Plant-derived ingredients are fully biodegradable and their production has a substantially lower ecological footprint compared to petrochemical-derived synthetic cosmetic ingredients.
- **Economic Accessibility:** Medicinal plants used in polyherbal hair oils are widely cultivated or wild-crafted and are significantly more affordable than pharmaceutical alternatives such as finasteride or specialized biologics.
- **Consumer Acceptance:** Growing consumer preference for 'clean,' 'natural,' and 'Ayurvedic' labels drives market adoption, particularly among consumers with sensitive skin or allergy concerns.

X. RECENT ADVANCES IN HERBAL HAIR OIL TECHNOLOGY

10.1 Nano-Herbal Formulations

The development of nanoencapsulation technologies has significantly expanded the therapeutic potential of herbal hair care formulations. Nanostructured lipid carriers (NLCs), solid lipid nanoparticles (SLNs), nanoemulsions, and phytosomes have been investigated as delivery systems for encapsulating bioactive phytoconstituents from Amla, Bhringraj, and Neem. Nanoencapsulation enhances bioavailability by improving follicular penetration, protects labile phytoconstituents from oxidative degradation, enables controlled release over extended periods, and reduces effective therapeutic doses.

10.2 Standardized Herbal Extracts

Standardized extracts defined by minimum assured concentrations of marker phytoconstituents (e.g.,

wedelolactone $\geq 0.2\%$ in Bhringraj, emblicanin A+B $\geq 5\%$ in Amla) are increasingly replacing crude powders in research-grade and commercial formulations. Standardization reduces inter-batch variability and enables dose-response relationship establishment critical for clinical evaluation.

10.3 Advanced Analytical Techniques

High-performance liquid chromatography (HPLC), ultra-high performance liquid chromatography coupled with mass spectrometry (UHPLC-MS), and nuclear magnetic resonance (NMR) spectroscopy are now routinely employed for phytochemical fingerprinting, authentication, and quality standardization of herbal hair oil raw materials and finished products. These techniques allow detection of adulterants, quantification of biomarkers, and elucidation of novel compounds.

10.4 Clinical and In Vivo Evidence

Randomized controlled trials have demonstrated statistically significant improvements in hair density, anagen-to-telogen ratio (assessed by trichoscopy), and patient-reported outcomes (hair fall, scalp itching, satisfaction) in subjects using polyherbal hair oil formulations compared to placebo mineral oil controls. While the clinical evidence base continues to expand, standardized outcome measures and longer follow-up periods are needed to establish definitive therapeutic claims.

XI. FUTURE SCOPE AND RESEARCH DIRECTIONS

Despite substantial progress, several areas merit further systematic investigation to fully realize the therapeutic potential of polyherbal hair oils and establish them as evidence-based cosmeceuticals:

8. **Rigorous Clinical Trials:** Large-scale, double-blind, placebo-controlled trials with validated trichoscopic outcome measures, standardized formulations, and adequate follow-up durations are essential for generating regulatory-grade clinical evidence.
9. **Mechanism Elucidation:** Molecular-level studies using transcriptomic, proteomic, and metabolomic approaches are needed to comprehensively characterize the signaling pathways modulated by polyherbal extracts in follicular cells.

10. Pharmacokinetic Studies: Investigation of follicular penetration profiles, percutaneous absorption rates, and bioavailability of key phytoconstituents from oil matrices will optimize formulation parameters.
11. Industrial Scale-Up and Standardization: Development of Good Manufacturing Practice (GMP)-compliant scale-up protocols and internationally harmonized quality standards is needed for consistent commercial production.
12. Novel Delivery Systems: Exploration of transferosomes, ethosomes, microemulsions, and carrier gel systems may further enhance follicular targeting and bioactive delivery efficiency.
13. Combination Therapy Studies: Evaluation of polyherbal hair oils as adjuvant therapy alongside standard-of-care treatments (minoxidil, ketoconazole shampoos) may demonstrate additive or synergistic clinical benefits.
14. Safety Pharmacology: Comprehensive dermal safety, sensitization, and phototoxicity assessments per OECD guidelines are needed to support regulatory submissions and international market authorization.

CONCLUSION

Polyherbal hair oils represent a scientifically validated and clinically relevant approach to the management of hair loss, scalp disorders, and impaired hair health. The convergent therapeutic activities of Amla, Bhringraj, Neem, Hibiscus, Fenugreek, and complementary botanical ingredients—encompassing antioxidant, anti-inflammatory, antimicrobial, hair growth-stimulating, and conditioning mechanisms—collectively provide a multi-targeted intervention that addresses the complex, multifactorial nature of hair disorders.

The integration of traditional ethnopharmacological knowledge with contemporary pharmaceutical sciences, advanced analytical characterization, and modern delivery technologies positions polyherbal hair oils as promising cosmeceutical candidates. However, the translation of *in vitro* and preclinical evidence into robust, regulatory-accepted clinical data requires sustained investment in standardized clinical research. With growing consumer preference

for natural, sustainable, and multi-functional cosmetic products, polyherbal hair oils are well-positioned to occupy a significant and expanding niche in the global herbal cosmetics market.

Future research efforts should prioritize clinical validation through randomized controlled trials, molecular mechanism characterization, pharmacokinetic optimization, and development of internationally harmonized quality standards to elevate polyherbal hair oil science from traditional practice to evidence-based cosmeceutical therapy.

Conflicts of Interest: The authors declare no conflicts of interest.

Funding: This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

REFERENCES

- [1] Kokate CK, Purohit AP, Gokhale SB. *Pharmacognosy*. 56th ed. Pune: Nirali Prakashan; 2021.
- [2] Trease GE, Evans WC. *Pharmacognosy*. 16th ed. London: Saunders Elsevier; 2009.
- [3] Sharma PP. *Cosmetics – Formulation, Manufacturing and Quality Control*. 4th ed. Delhi: Vandana Publications; 2010.
- [4] World Health Organization. *WHO Guidelines on Good Herbal Processing Practices (GHPP) for Herbal Medicines*. Geneva: WHO; 2018.
- [5] Adhirajan N, Ravi Kumar T, Shanmugasundaram N, Babu M. *In vivo* and *in vitro* evaluation of hair growth potential of *Hibiscus rosa-sinensis* Linn. *J Ethnopharmacol*. 2003;88(2-3):235–239.
- [6] Datta K, Singh AT, Mukherjee A, Bhat B, Ramesh B, Burman AC. *Eclipta alba* extract with potential for hair growth promoting activity. *J Ethnopharmacol*. 2009;124(3):450–456.
- [7] Nayak BS, Raju SS, Chalapathi Rao AV. Wound healing activity of *Persea americana* (avocado) fruit: a preclinical study on rats. *J Wound Care*. 2008;17(3):123–126.
- [8] Rele AS, Mohile RB. Effect of mineral oil, sunflower oil, and coconut oil on prevention of

- hair damage. *J Cosmet Sci.* 2003;54(2):175–192.
- [9] Pandit S, Bhaskaran D, Prasad LK, Priya K. Physicochemical and phytochemical evaluation of polyherbal hair oil. *Asian J Pharm Clin Res.* 2017;10(3):152–157.
- [10] Joshi SG. *Medicinal Plants.* New Delhi: Oxford & IBH Publishing; 2000.
- [11] Mukherjee PK, Maity N, Nema NK, Sarkar BK. Bioactive compounds from natural resources against skin aging. *Phytomedicine.* 2011;19(1):64–73.
- [12] Patel S, Sharma V, Chauhan NS, Dixit VK. Hair growth: Focus on herbal therapeutic agents. *Curr Drug Discov Technol.* 2015;12(1):21–42.
- [13] Semalty M, Semalty A, Joshi GP, Rawat MSM. Hair growth and rejuvenation: An overview. *J Dermatol Treat.* 2011;22(3):123–132.
- [14] Uslu M, Senel E, Bayraktar N. Effect of herbal extracts on the 5-alpha reductase inhibition: an in vitro study. *Phytother Res.* 2012;26(11):1654–1659.
- [15] Rajput RJ. Influence of nutrition, food supplements and lifestyle in hair disorders. *Int J Trichol.* 2017;9(2):58–67.