

A Traditional Potassium Permanganate–Tamarind Pulp Preparation Used for Scorpion Stings in Rural Maharashtra, India: An Ethnopharmacological Observation and Testable Mechanistic Hypothesis

S. V. Deshpande

Dr. D. Y. Patil Institute of Pharmacy, Akurdi, Pune, Maharashtra, India

Published October 2026

Abstract

Traditional practices used for scorpion stings remain part of community experience in some rural settings, but they require careful documentation and critical evaluation. This article describes a practice reported and observed by the author in rural western Maharashtra, India, in which a small quantity of potassium permanganate (KMnO_4) crystals is placed on moist pulp from ripe tamarind (*Tamarindus indica* L.), and the preparation is applied to the sting site. The observations were accumulated informally over several years and were not collected through a prospective study, structured interviews, standardized pain scoring, or systematic species identification. Some episodes were reported to be followed by a reduction in local pain, while others reportedly showed little or no apparent benefit. These observations cannot establish efficacy or causation. The combination raises a testable chemical question because permanganate is a strong oxidizing agent, whereas tamarind pulp is an acidic, chemically complex matrix. Oxidative modification of accessible venom components is conceivable in principle, but topical contact with skin cannot be assumed to reach venom injected into tissue or to neutralize it. Alternative explanations—including counter-irritation, expectation effects, and the natural course of pain—must also be considered. This paper documents the practice as a hypothesis-generating observation and outlines a research pathway involving preparation characterization, controlled in vitro testing, peptide analysis, functional assays, and safety assessment. The preparation is not recommended as treatment; suspected scorpion envenomation requires prompt medical assessment.

Keywords: scorpion sting; traditional knowledge; ethnopharmacology; potassium permanganate; *Tamarindus indica*; venom peptides; rural Maharashtra

1 Introduction

Scorpion stings are a relevant health concern in several tropical and subtropical regions. Clinical effects vary by species, the amount of venom delivered, the site of the sting, and the individual affected. Symptoms may range from severe localized pain to systemic manifestations requiring urgent medical care. In India, the Indian red scorpion (*Mesobuthus tamulus*) is among the medically important species, including in parts of Maharashtra. Scorpion venoms are complex mixtures containing peptides and other components; several neurotoxins affect voltage-gated ion channels and can alter neuronal excitability. Communities may use local remedies in response to the immediate pain of a sting, particularly where access to healthcare is delayed. Documenting such practices can contribute to the study of traditional knowledge, but documentation must be distinguished from evidence of safety or efficacy. A practice may be culturally established without having been clinically evaluated,

and a reported temporal association between application and pain reduction does not demonstrate that the preparation caused the change.

This paper describes a potassium permanganate–tamarind preparation encountered in rural western Maharashtra. It does not report a clinical trial, a formal ethnobotanical survey, or laboratory findings. Its purpose is narrower: to document the practice as an informal field observation, identify the limits of the available information, and formulate competing hypotheses that could be investigated without endorsing the preparation for human use.

2 Basis and limits of the observation

The account is based on the author's informal observations and community reports over several years in rural parts of Satara, Sangli, and Solapur districts of Maharashtra. The practice was encountered in agricultural and village settings, particularly during warmer periods and the early monsoon. No prospective protocol was used. There was no structured questionnaire, predefined case definition, systematic recruitment, independent verification of each episode, or standardized clinical follow-up. The number of distinct individuals and episodes cannot be reliably reconstructed from the available notes and is therefore not quantified here.

In the practice as described, moist pulp from a ripe tamarind fruit is used as a matrix, and a small, unmeasured quantity of potassium permanganate crystals with some water drops is placed on the pulp. The resulting material is rubbed or placed at the reported sting site. Community accounts and the author's observations described a burning or heating sensation and, in some episodes, a subsequent reduction in local pain within minutes. Other episodes reportedly showed little or no apparent relief. The study did not consistently record the amount of KMnO_4 , the interval between sting and application, duration of contact, scorpion species, venom dose, co-interventions, or objective outcomes. Whilst doctors treat with local anesthetic injections for immediate pain relief, after some time the pain persists, as observed.

These limitations are central rather than incidental. Pain can change over time, perceived improvement can be influenced by expectation, and the identity and severity of the sting may vary. The observation cannot determine how often improvement occurred, whether the practice caused it, or whether adverse effects were overlooked. Accordingly, the description should be interpreted as an account of a reported traditional practice, not as a clinical case series or evidence of therapeutic benefit.

3 Scientific context and testable hypotheses

3.1 Scorpion venom and peptide structure

Many scorpion neurotoxins are compact peptides stabilized by disulfide bonds and interact with voltage-gated sodium or potassium channels. Their biological effects depend on molecular structure and interactions with their targets; this makes toxin structure a legitimate topic for mechanistic research, but it does not by itself show that a topical oxidant can inactivate venom in living tissue.

3.2 Potassium permanganate and the tamarind matrix

Potassium permanganate is a strong oxidizing agent. Under appropriate experimental conditions, it can react with susceptible organic molecules. Proteins and peptides contain residues that may be

oxidatively modified, although the products and extent of reaction depend on the chemical environment. Ripe tamarind pulp contains organic acids, sugars, pectin, and phenolic constituents, and its composition may vary with cultivar, ripeness, processing, and storage. These components may alter the redox chemistry of a preparation rather than simply acting as an inert carrier.

The central hypothesis is that the combined preparation might chemically modify some accessible venom components under defined conditions. This is only a hypothesis. It is not established that the preparation reaches venom deposited beneath the skin, that any chemical modification would reduce toxin activity, or that such a reaction could account for a perceived change in pain. The acidic and chemically complex pulp may also consume the oxidant, and reactions at the skin surface cannot be assumed to reproduce conditions in tissue.

3.3 Competing explanations

At least four explanations should be considered: (i) a direct chemical interaction with a fraction of accessible venom, if such contact occurs; (ii) a local sensory or counter-irritant effect associated with burning or other skin sensations; (iii) expectation-related changes in reported pain; and (iv) the natural evolution of pain after a sting. More than one process could occur, and the current observations cannot discriminate among them. The possibility of chemical irritation or burns is also a safety concern, not evidence of therapeutic action.

3.4 Proposed research pathway

A staged programme could test the hypothesis without exposing people to an unvalidated preparation. First, the preparation itself should be characterized using reproducible, laboratory-defined quantities rather than the unmeasured amounts used in the traditional practice. Relevant measurements may include pH, oxidation–reduction potential, visible/UV spectroscopy, reaction kinetics, and appropriate assessment of manganese-containing products. Preliminary safety and chemical compatibility work should precede any venom experiments.

Second, controlled in vitro experiments could compare venom alone with venom exposed separately to KMnO_4 , a standardized tamarind preparation, and their combination, alongside suitable reagent and matrix controls. The work should use justified concentrations and exposure times established by preliminary analytical studies; it should not attempt to recreate an uncontrolled human application. LC–MS/MS could assess whether reproducible peptide modifications occur. Orthogonal structural methods may be useful where appropriate, but structural change alone is not evidence of neutralization.

Third, any detected chemical changes should be followed by validated functional assays relevant to the venom and toxin under study. Only a reproducible reduction in biological activity, with adequate controls, would support further investigation. Subsequent safety studies would need to assess local tissue injury and systemic toxicity under institutional oversight. Human studies should not be contemplated unless substantial preclinical evidence and the necessary ethical and regulatory approvals are in place.

An ethnographic follow-up, if undertaken, should be separately designed. It should use a transparent sampling approach, informed consent, appropriate institutional ethics review, and a structured interview guide that records how the practice is learned, when and where it is used, perceived outcomes, failures, adverse effects, and whether medical care was sought. It should not encourage

participants to apply the preparation or delay care. Any report should distinguish firsthand observation from secondhand community testimony.

3.5 Safety and public-health considerations

Potassium permanganate is a strong oxidizer and can irritate or injure tissue; concentrated or prolonged contact can cause chemical burns. Because the traditional preparation is unstandardized and has not been assessed for safety or efficacy, this article does not recommend applying it to a sting. A scorpion sting can produce systemic complications, and local pain intensity alone does not reliably exclude serious envenomation. Prompt medical assessment and evidence-based management should take precedence. No traditional preparation should delay transport, monitoring, analgesia, or other treatment indicated by a qualified clinician.

4 Conclusion

A potassium permanganate–tamarind preparation has been reported as a traditional response to scorpion stings in rural western Maharashtra. The observation is worth documenting because it raises questions about local knowledge, the chemistry of a mixed oxidizing preparation, and the interpretation of perceived pain changes. However, the available information does not establish efficacy, mechanism, or safety. The most defensible next step is rigorous documentation of the practice and controlled laboratory investigation of its chemistry and any functional effects, while maintaining a clear warning against its use as a substitute for medical care.

References

- [1] Isbister GK, Bawaskar HS. Scorpion envenomation. *N Engl J Med*. 2014;371:457–463.
- [2] Rodríguez de la Vega RC, Possani LD. Current views on scorpion toxins specific for K⁺ channels. *Toxicon*. 2004;43:865–875.
- [3] Ortiz E, Gurrola GB, Schwartz EF, Possani LD. Scorpion venom components as potential candidates for drug development. *Toxicon*. 2015;93:125–135.
- [4] Petricevich VL. Scorpion venom and the inflammatory response. *Mediators Inflamm*. 2010;2010:903295.
- [5] Bawaskar HS, Bawaskar PH. Scorpion sting: update. *J Assoc Physicians India*. 2012;60:46–55.
- [6] Pucca MB, Cerni FA, Janke R, et al. History of envenoming therapy and current perspectives. *Front Immunol*. 2019;10:1598.
- [7] De Caluwé E, Halamová K, Van Damme P. Tamarindus indica L. – a review of traditional uses, phytochemistry and pharmacology. *Afr Focus*. 2010;23:53–83.
- [8] Sudjaroen Y, Haubner R, Würtele G, et al. Isolation and antioxidant activity of phenolic compounds from tamarind. *Food Chem Toxicol*. 2005;43:1673–1682.
- [9] Patwardhan B, Vaidya ADB, Chorghade M. Reverse pharmacology and systems approaches for drug discovery. *J Ethnopharmacol*. 2008;115:115–118.
- [10] Barth A. Infrared spectroscopy of proteins. *Biochim Biophys Acta*. 2007;1767:1073–1101.
- [11] Kelly SM, Jess TJ, Price NC. How to study protein structure by circular dichroism. *Biochim Biophys Acta*. 2005;1751:119–139.
- [12] Ediriweera ERHSS, Ratnasooriya WD, Kamathewatta KWM. *Ethnomedicinal uses in scorpion poisoning (Gonusu Visha) in Sri Lankan traditional medicine*. Int J Ayurveda Pharma Res. 2018.